

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 5/06, 5/08, 5/10, A61K 35/12, 48/00		A2	(11) International Publication Number: WO 99/64565
			(43) International Publication Date: 16 December 1999 (16.12.99)
(21) International Application Number: PCT/US99/12847		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 8 June 1999 (08.06.99)		Published <i>Without international search report and to be republished upon receipt of that report.</i>	
(30) Priority Data:			
60/088,431 8 June 1998 (08.06.98) US			
60/099,233 4 September 1998 (04.09.98) US			
(71) Applicant: OSIRIS THERAPEUTICS, INC. [US/US]; 2001 Aliceanna Street, Baltimore, MD 21231-2001 (US).			
(72) Inventors: THIEDE, Mark, A.; 319 Pond View Court, Forest Hill, MD 21050 (US). MBALAVIELE, Gabriel; Apartment 1G, 8488 Greystone Lane, Columbia, MD 21045 (US).			
(74) Agents: OLSTEIN, Elliot, M. et al.; Carella, Byrne, Bain, Gilfillan, Cecchi, Stewart & Olstein, 6 Becker Farm Road, Roseland, NJ 07068 (US).			
(54) Title: REGULATION OF HEMATOPOIETIC STEM CELL DIFFERENTIATION BY THE USE OF HUMAN MESENCHYMAL STEM CELLS			
(57) Abstract			
<p>The invention relates to the induction of hematopoietic stem cells to differentiate into osteoclasts by culturing hematopoietic stem cells with human mesenchymal stem cells, and, in a preferred embodiment, using no exogenous cytokines. Differentiation of the mesenchymal stem cells into osteoblasts inhibited the differentiation of hematopoietic stem cells into osteoclasts. In addition, hematopoietic stem cells can be genetically engineered to carry genes of interest particularly for the expression of physiologically active proteins. In the presence of mesenchymal stem cells, the transduced cells carry the new genetic material and express gene products that can be used to modulate bone resorption.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakhstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

5

10 **REGULATION OF HEMATOPOIETIC STEM CELL DIFFERENTIATION
BY THE USE OF HUMAN MESENCHYMAL STEM CELLS**

 The present invention relates to hematopoietic stem cells and more particularly to a process and composition for differentiating human hematopoietic stem cells into a committed osteoclast lineage in the absence of exogenous
15 osteoclastogenesis-inducing factors. The present invention further relates to genetically modified hematopoietic stem cells in the presence of human mesenchymal stem cells, such that when the hematopoietic stem cells differentiate into osteoclasts, the osteoclasts are able to express the product of the transduced
20 gene.

Background of the Invention

 Osteoclasts are terminally differentiated hematopoietic cells responsible for physiological bone resorption. Dysfunction in osteoclast differentiation and/or
25 activity is the cause of a variety of human metabolic bone diseases including osteoporosis (for review see Teitelbaum et al., 1995). Osteoporosis, or progressive bone loss, is attributed to a shift in the critical balance of osteoclast and osteoblast activities responsible for proper skeletal integrity. Osteoblasts, which build bone, are one terminally differentiated lineage descending from mesenchymal stem cells.

30

 Hematopoietic progenitors in bone marrow are driven into the osteoclast cell lineage under the influence of local regulatory factors. Diverse cell types including stromal cells, which arise from mesenchymal stem cells (MSCs), are present in bone marrow. Stromal cells have been shown to produce factors that
35 regulate osteoclast physiology (for review see Mundy et al., 1995). Histological

pictures of bone show close vicinity between marrow stromal cells and hematopoietic cells. Further, the relationship between the differentiated status of bone marrow stromal cells and their potential to regulate neighboring osteoclast progenitors is poorly documented. The importance of heterotypic cell-to-cell interactions mediated by cadherin-6 between stromal cells and osteoclast progenitors using a mouse model of osteoclastogenesis has been reported (Mbalaviele et al., 1998). Several studies demonstrated that osteoclast differentiation required the presence of stromal/osteoblast cells (Takahashi et al., 1988; Mbalaviele et al., 1995). Recent studies using cells derived from rodent marrow have demonstrated that osteoprotegerin ligand (OPGL) expressed on the membrane of stromal cells is a key factor of osteoclastogenesis (Lacey et al. 1998; Kong et al. 1999). However, it has also been reported that human osteoclast cells differentiated from progenitor cells in the presence of cocktails of cytokines and that stromal cells were not necessary (Matayoshi et al., 1996). The formation of human osteoclast cells (Ocl) displaying bone-resorbing activity has been reported (Takahashi et al., 1995; James et al., 1996; Sarma et al., 1996; Quinn et al., 1997), however, critical events which take place during osteoclastogenesis are unclear.

Bone is a composite matrix consisting of both inorganic and organic elements. The organic phase contains proteins, while the inorganic component contains calcium salts. It is known that during osteoclastic bone resorption, dissolution of the inorganic phase precedes that of the organic phase and that high concentrations of calcium (>40mM) are released locally from bone matrix. In turn, increased calcium concentrations have a negative feedback on osteoclast function, a regulatory mechanism to prevent excessive bone resorption. Calcium is known to regulate the expression of several genes. Recently, a calcium response element that enhances human keratin-1 gene expression after exposure of keratinocytes to extracellular calcium (0.6 mM) has been characterized.

Osteoclasts are not easily isolated due to their low cell number *in vivo*, their fragility and tendency to adhere to other bone cells. Thus, there is a need for a method of efficiently producing quantities of osteoclasts to facilitate, for example, the design of effective therapeutics aimed at preventing abnormal osteolysis.

Accordingly, it is an object of the present invention to provide an efficient method of producing osteoclasts *in vitro*.

It is a further object of the invention to provide a method to produce and
5 maintain osteoclasts *in vitro* without the addition of exogenous growth factors in numbers sufficient to provide an ongoing source of osteoclasts, for example, for administering osteoclasts to an individual in need thereof.

Accordingly, it is an object of the present invention to regulate
10 hematopoietic stem cell differentiation towards osteoclast cells in the absence of exogenous growth factors, cytokines and hormones.

It is a still further object of the present invention to provide a method of
producing osteoclasts from genetically modified hematopoietic cells such that
15 when the hematopoietic cells differentiate into osteoclasts, the osteoclasts express the gene product of interest.

Summary of the Invention

The present inventors have discovered that mesenchymal stem cells support
20 osteoclast cell differentiation of hematopoietic stem cells.

Thus in one aspect of the present invention, there is provided a method for inducing the differentiation of hematopoietic progenitor cells *in vitro* into osteoclast cells, comprising co-culturing the hematopoietic cells with human
25 mesenchymal stem cells.

When hematopoietic progenitor cells were co-cultured with mesenchymal stem cells, the hematopoietic progenitor cells differentiated towards osteoclast cells which were identified by the expression of markers characteristic of osteoclasts
30 including multinuclearity, calcitonin and vitronectin receptors, and more importantly, bone resorbing activity.

It has further been discovered that the differentiation of the hematopoietic stem cells occurred in the absence of exogenous (added) growth factors, cytokines

and hormones known to be required for inducing differentiation of hematopoietic stem cells.

Accordingly, one aspect of the present invention provides a method of
5 producing osteoclasts *in vitro* comprising co-culturing hematopoietic stem cells with human mesenchymal stem cells. In a preferred embodiment, the cells are co-cultured in the same culture vessel such that physical cell-to-cell interactions occur between the hematopoietic progenitor cells and the mesenchymal stem cells.

10 It has also been discovered that when hematopoietic stem cells, which have been modified to carry exogenous genetic material of interest, are co-cultured with mesenchymal stem cells, the transduced hematopoietic stem cells differentiated into osteoclasts that also carried the new genetic material. These transduced
15 osteoclast cells are able to express the exogenous gene product. Transduced osteoclast progenitors and the osteoclasts differentiated therefrom can be used for applications where treatment using such modified osteoclasts is beneficial, for example, in the alleviation of the effects of osteoporosis.

Accordingly, the present invention provides a method of obtaining
20 genetically modified osteoclasts, comprising transducing hematopoietic progenitor cells with exogenous genetic material and placing the transduced hematopoietic cells under conditions suitable for differentiation of the hematopoietic stem cells into osteoclast cells which contain the exogenous genetic material.

25 In one embodiment, the method of producing osteoclasts comprises co-culturing transduced hematopoietic stem cells with mesenchymal stem cells such that after differentiation of the hematopoietic stem cells into osteoclasts, the osteoclasts also contain the exogenous genetic material.

30 The invention further relates to hematopoietic stem cells that are transduced with a polynucleotide that encodes a product that downregulates osteoclast differentiation and/or activity, and the use thereof.

In a particularly preferred embodiment, the expressed gene product downregulates osteoclast differentiation and/or activity. Hematopoietic stem cells are transduced with a retroviral vector containing a tandem of a calcium response element, tartrate resistant acid phosphatase (TRAP) promoter (specifically expressed in osteoclasts) and a nucleic acid sequence encoding an anti-osteoclast activity factor (for example, an antisense sequence to M-CSF, IL1 or IL6; an inhibitor such as an intracellular antibody to IL6, to M-CSF, to GM-CSF, to IL1, or to Leukemia Inhibitory Factor; or, an apoptotic factor). Expression of this sequence is triggered by the local increase in calcium concentration due to osteoclastic bone resorption.

It has further been found that while mesenchymal stem cells positively regulated the differentiation of CD34⁺ progenitor cells towards osteoclast cells, MSCs that have begun differentiation into osteoblastic cells inhibited this process. In the presence of osteogenic mesenchymal stem cells, the levels of OPGL expression decreased and expression of osteoprotegerin (OPG) were found to increase. Thus, it is also contemplated that culturing hematopoietic stem cells in the presence of differentiated mesenchymal stem cells may serve to downregulate osteoclast differentiation and/or activity. Accordingly, the co-cultured cell populations of the invention may be used as part of a treatment to alleviate the symptoms of osteoporosis.

Brief Description of the Drawings

Figure 1A shows TRAP staining of CD34⁺ cells cultured in the absence of mesenchymal stem cells (MSCs).

Figure 1B shows TRAP staining of CD34⁺ cells cultured in the presence of MSCs. Many multinucleated TRAP⁺ cells (TRAP+MNCs) (arrow) as well as mononuclear TRAP⁺ cells were formed in the presence of MSCs which appeared as a confluent layer of spindle-shaped cells (arrowhead). Small arrows and star indicate clusters of nuclei.

Figure 1C shows staining of CD34⁺ cells cultured in the presence of human skin fibroblasts (SK1087 cells; arrowhead). No TRAP+MNCs were formed.

Figure 1D shows staining of CD34⁺ cells cultured in the presence of human kidney cells (293T; arrowhead). TRAP+MNCs were not formed.

Figure 2A shows immunocytochemical staining of co-cultures of CD34⁺ and MSCs with negative-control antibody.

Figure 2B shows immunocytochemical staining of co-cultures of CD34⁺ and MSCs with antibody recognizing vitronectin receptor.

Figure 3 shows RT-PCR of RNA purified from co-cultures of MSCs and CD34⁺ cells (lanes 2-3) or cultures of MSCs without CD34⁺ cells (lanes 4-5). Calcitonin receptors (lanes 2 and 4) and TRAP (lanes 3 and 5) were only detected in samples from co-cultures of MSCs and CD34⁺ cells. The conserved and unspliced region of the calcitonin receptor family was chosen to design primers. Lane 1, 1 kb DNA ladder; bottom, β 2-microglobulin.

Figure 4A shows pit formation in co-cultures of CD34⁺ cells and MSCs (arrows).

Figure 4B shows pit formation in cultures of CD34⁺ cells without MSCs.

Slices were stained with toluidine blue and visualized under light microscopy. X200 magnification.

Figure 4C shows quantitation of pits in co-cultures of CD34⁺ cells and MSCs (closed bar) or in cultures of CD34⁺ cells without MSCs (open bar). * $p < 0.01$ versus cultures of CD34⁺ cells without MSCs.

Figure 5A shows the effects of cell-to-cell interactions in osteoclast cell formation. CD34⁺ cells and MSCs were co-cultured in the same compartments (contact co-cultures, closed bar) or separated by means of 0.45 μ m filters (non-contact co-cultures): MSCs on top and CD34⁺ cells on the bottom, open bar; MSCs on the bottom and CD34⁺ cells on top, dashed bar). * $p < 0.01$ versus non-contact co-cultures.

Figure 5B shows quantitation of secreted factors (IL-6, IL-11, LIF) in co-cultures of CD34⁺ cells with MSCs. The levels of IL-6 and LIF but not IL-11 were about 10 times higher in co-cultures of MSCs and CD34⁺ cells compared to MSC cultures without CD34⁺ cells

Figure 6A and 6B show inhibition of TRAP+MNCs by osteogenic MSCs. Osteogenic differentiation of MSCs was induced by treatment with osteogenic supplement (OS) for 2, 3, 4, 5, 10 or 13 days prior to adding CD34+ cells. * $p < 0.01$ versus co-cultures of OS-treated MSCs and CD34+ cells.

5 Figure 6C shows that co-cultures of MSCs and CD34+ cells produced higher levels of IL-6, IL-11 and LIF than co-cultures of osteogenic MSCs and CD34+ cells.

Figure 6D shows that osteogenic differentiation of MSCs is associated with a decrease in OPGL mRNA expression (left panel) and an increase in OPG mRNA expression (right panel).

10

Figure 7A shows flow cytometry analysis of CD34⁺ cells transduced with control vector.

Figure 7B shows flow cytometry analysis of CD34⁺ cells transduced with EGFP expression vector. Approximately 30% of CD34⁺ cells expressed EGFP.

15

Figure 7C shows co-cultures of MSCs with CD34⁺ cells transduced with EGFP expression vector. EGFP was expressed by osteoclasts as well as by mononuclear cells.

Figure 7D shows co-cultures of MSCs with CD34⁺ cells transduced with EGFP expression vector stained for TRAP. Fig. 7D is the light view of Fig. 7C.

20

Figure 8A shows the expression of vitronectin receptor ($\alpha_v\beta_3$) detected by ELISA. $\alpha_v\beta_3$ antibody, wells on left and closed bar; control IgG, wells on right and open bar.

Figure 8B shows the time-course expression of $\alpha_v\beta_3$. Co-cultures of CD34+ cells and MSCs (closed bars) or MSCs without CD34+ cells (open bars).

25

Figure 9A shows a strategy for direct regulation of osteoclast activity.

Figure 9B shows that the addition of calcium to cultured cells transfected with a calcium-inducible expression system induced expression of the reporter gene.

30

Figure 10 shows the effect of serum in culture medium on production of osteoclasts in co-cultures of MSCs and CD34+ cells.

Figure 10A shows osteoclast numbers in co-cultures of CD34+ cells and MSCs in DMEM-low glucose with 10% FBS, open bar; and in DMEM-high glucose without serum, solid bar.

Figure 10B show co-cultures of 293 cells and CD34+ cells in DMEM-high glucose with 10% FBS, solid bar; and in DMEM-high glucose without serum, dashed bar.

Detailed Description of the Invention

The present invention relates generally to the use of human mesenchymal stem cells to induce the differentiation of human CD34+ cells into osteoclast cells and compositions comprising human CD34+ cells and human mesenchymal stem cells. More particularly, the inventors found that human mesenchymal stem cells cultured in association with CD34+ cells are useful for inducing the differentiation of CD34+ cells into osteoclasts. Thus the CD34+ cells can be maintained and utilized as a source of osteoclasts.

In order to obtain subject human mesenchymal stem cells for the methods described herein, mesenchymal stem cells can be recovered from bone marrow or other mesenchymal stem cell source. Bone marrow cells may be obtained from iliac crest, femora, tibiae, spine, rib or other medullary spaces. Other sources of human mesenchymal stem cells include embryonic yolk sac, placenta, umbilical cord, fetal and adolescent skin, and blood. The presence of mesenchymal stem cells in the culture colonies may be verified by specific cell surface markers which are identified with unique monoclonal antibodies, see, e.g., U.S. Patent No. 5,486,359. These isolated mesenchymal cell population display epitopic characteristics associated only with mesenchymal stem cells, have the ability to regenerate in culture without differentiation and have the ability to differentiate into specific mesenchymal lineages when either induced in vitro or in vivo at the site of damaged tissue. Human mesenchymal stem cells are known to differentiate under the appropriate conditions into multiple cell lineages, including osteoblasts, adipocytes and chondrocytes (for review see Caplan et al., 1997).

Accordingly, any process that is useful to recover mesenchymal stem cells from human tissue may be utilized to obtain a population of cells comprising enriched mesenchymal stem cells. In one aspect, the method of isolating human mesenchymal stem cells comprises the steps of providing a tissue specimen
5 containing mesenchymal stem cells, preferably bone marrow; isolating the mesenchymal stem cells from the specimen, for example, by density centrifugation; adding the isolated cells to a medium which contains factors that stimulate mesenchymal stem cell growth without differentiation, and allows for the selective adherence of only the mesenchymal stem cells to a substrate surface in culture;
10 culturing the specimen-medium mixture; and removing the non-adherent matter from the substrate surface.

In a further aspect of the present invention, any process that is useful to recover hematopoietic stem cells from human tissue may be utilized to result in a
15 population of cells comprised mostly of hematopoietic cells. Stem cells can be recovered from various types of tissue such as bone marrow and blood, including peripheral blood. The human hematopoietic stem cells can be collected from bone marrow aspirates or peripheral blood and isolated using commercially available antibodies which bind to hematopoietic stem cell surface antigens, e.g. CD34,
20 using methods known to those of skill in the art, see e.g. U.S. Pat. No. 4,714,680. Alternatively, the hematopoietic stem cells can be recovered by depletion, i.e., by selecting out cells which express markers found on other cells. The antibodies used for these procedures may be conjugated to magnetic beads and immunogenic procedures may be utilized to recover the desired cell type.

25

The human mesenchymal stem cells and the hematopoietic stem cells are co-cultured under appropriate culture conditions such that the mesenchymal stem cells adhere to a substrate surface and form a monolayer. The mesenchymal stem cells are plated at a density in a range of from about 3×10^3 to about 5×10^3 cells
30 per cm^2 . The CD34+ cells are preferably at a cell density of approximately 5×10^4 cells per cm^2 .

According to the method of the present invention, the isolated mesenchymal stem cells and the isolated hematopoietic progenitor cells, preferably CD34+ cells, are

each culture expanded in appropriate media, i.e., cultured by methods using conditions that are apparent to those of skill in the art which favor cell growth and production of homogeneous cell populations.

5 The two cell populations are then co-cultured in a medium that promotes the growth of the human mesenchymal stem cells and does not adversely affect the maintenance of the hematopoietic stem cells. The media should also sustain the CD34+ cell population and support the induction of the differentiation of CD34+ cells into osteoclast cells. Suitable media are described for example in U.S. Pat. No.
10 5,486,359.

 In a preferred embodiment, the human mesenchymal stem cells and the hematopoietic stem cells are co-cultured in Dulbecco's Modified Eagles Medium-Low Glucose (DMEM-LG #11885 Life Technologies, Gaithersburg, MD). The medium
15 preferably contains 10% FBS. In a particularly preferred embodiment, the human mesenchymal stem cells and the hematopoietic stem cells are co-cultured in Dulbecco's Modified Eagles Medium-High Glucose (DMEM-HG # 11995 Life Technologies, Gaithersburg, MD) without FBS.

20 The hematopoietic stem cells produced according to the methods described herein can be used to provide a reliable and constant source of osteoclast cells. These osteoclasts could be used in the determination of the effects of osteotropic factors on the development and treatment of osteoporosis. It is contemplated that a ready supply of osteoclasts would be useful for screening assays for small molecule
25 drug development, e.g., based on expression of integrin $\alpha_v\beta_3$ cell surface receptor by osteoclasts. It is believed that these cells, which have not been treated with growth factors *in vitro*, will have characteristics more like osteoclasts found *in vivo* than have heretofore been obtained.

30 After differentiation of the hematopoietic progenitor cells into osteoclast cells as described herein, the mixture of hematopoietic progenitor cells, osteoclasts and mesenchymal stem cells may be separated to obtain a population of cells largely consisting of the osteoclast cells. This may be accomplished by positive

and/or negative selection of hematopoietic cells using antibodies to identify hematopoietic cell surface markers or other cell-type markers.

Another aspect of the present invention relates to the introduction of foreign
5 genes into the hematopoietic progenitor cells such that progeny of the cells, the osteoclasts, carry the new genetic material.

Thus, in accordance with this aspect of the invention, the hematopoietic progenitor cells can be modified with genetic material of interest. The modified
10 CD34+ cells can then be co-cultured *in vitro* with mesenchymal stem cells and induced to differentiate into osteoclast cells. The osteoclast cells are able to express the product of the gene expression or secrete the expression product. These modified cells can then be administered to a target tissue, e.g. bone marrow, where the expressed product will have a beneficial effect.

15

In a further embodiment, the transduced hematopoietic progenitor cells can be induced *in vivo* to differentiate into osteoclasts that will express the gene product. For example, the transduced hematopoietic progenitor cells in combination with mesenchymal stem cells may be administered to induce
20 production of osteoclasts having the transduced gene. The cells may be administered in admixture with each other or separately and may be delivered to a targeted area. Alternatively, the hematopoietic progenitor cells may be used alone and caused to differentiate *in vivo* by mesenchymal stem cells present *in vivo*.

25

Thus, genes can be introduced into cells which are then returned to the autologous donor or an allogeneic recipient where the expression of the gene will have a therapeutic effect. For example, osteoclasts may be genetically engineered to have reduced activity *in vivo*. Appropriate genes would include those that play a role in the regulation of osteoporosis, in areas such as serum calcium
30 responsiveness, estrogen secretion and bone resorption.

The hematopoietic stem cells may be genetically modified (transduced or transformed or transfected) in the presence of the human mesenchymal stem cells, wherein the mesenchymal stem cells increase the efficiency of gene transduction of

the hematopoietic stem cells. Alternatively, the hematopoietic stem cells may be transduced in the absence of the human mesenchymal stem cells.

5 The hematopoietic stem cells may be genetically modified by incorporation of genetic material into the cells, for example using recombinant expression vectors.

As used herein "recombinant expression vector" refers to a transcriptional unit comprising an assembly of (1) a genetic element or elements having a regulatory role in gene expression, for example, promoters or enhancers, (2) a structural or coding
10 sequence which is transcribed into mRNA and translated into protein, and (3) appropriate transcription initiation and termination sequences. Structural units intended for use in eukaryotic expression systems preferably include a leader sequence enabling extracellular secretion of translated protein by a host cell. Alternatively, where recombinant protein is expressed without a leader or transport
15 sequence, it may include an N-terminal methionine residue. This residue may or may not be subsequently cleaved from the expressed recombinant protein to provide a final product.

The human hematopoietic progenitor cells thus may have stably integrated a
20 recombinant transcriptional unit into chromosomal DNA or carry the recombinant transcriptional unit as a component of a resident plasmid. Cells may be engineered with a polynucleotide (DNA or RNA) encoding a polypeptide *ex vivo*, for example. Cells may be engineered by procedures known in the art by use of a retroviral particle containing RNA encoding a polypeptide.

25 Retroviruses from which the retroviral plasmid vectors hereinabove mentioned may be derived include, but are not limited to, Moloney Murine Leukemia Virus, spleen necrosis virus, retroviruses such as Rous Sarcoma Virus, Harvey Sarcoma Virus, avian leukosis virus, gibbon ape leukemia virus, human
30 immunodeficiency virus, adenovirus, Myeloproliferative Sarcoma Virus, and mammary tumor virus. In one embodiment, the retroviral plasmid vector is MGIN, derived from murine embryonic stem cells. Generally regarding retroviral mediated gene transfer, see McLachlin et al.(1990).

The nucleic acid sequence encoding the polypeptide is under the control of a suitable promoter. Suitable promoters which may be employed include, but are not limited to, TRAP promoter, adenoviral promoters, such as the adenoviral major late promoter; the cytomegalovirus (CMV) promoter; the respiratory syncytial virus (RSV) promoter; inducible promoters, such as the MMT promoter, the metallothionein promoter; heat shock promoters; the albumin promoter; the ApoAI promoter; human globin promoters; viral thymidine kinase promoters, such as the Herpes Simplex thymidine kinase promoter; retroviral LTRs; ITRs; the β -actin promoter; and human growth hormone promoters. The promoter also may be the native promoter that controls the gene encoding the polypeptide. These vectors also make it possible to regulate the production of the polypeptide by the engineered progenitor cells. For example, for purposes of the present invention, the vector can contain a calcium response element. The selection of a suitable promoter will be apparent to those skilled in the art.

It is also possible to use vehicles other than retroviruses to genetically engineer or modify the hematopoietic stem cells. Genetic information of interest can be introduced by means of any virus which can express the new genetic material in such cells. For example, SV40, herpes virus, adenovirus and human papillomavirus can be used for this purpose. Other methods can also be used for introducing cloned eukaryotic DNAs into cultured mammalian cells, for example, the genetic material to be transferred to stem cells may be in the form of viral nucleic acids.

In addition, the expression vectors may contain one or more selectable marker genes to provide a phenotypic trait for selection of transformed cells such as dihydrofolate reductase or neomycin resistance.

The hematopoietic progenitor cells may be transfected through other means known in the art. Such means include, but are not limited to transfection mediated by calcium phosphate or DEAE-dextran; transfection mediated by the polycation Polybrene; protoplast fusion; electroporation; liposomes, either through encapsulation of DNA or RNA within liposomes, followed by fusion of the liposomes with the cell

membrane or, DNA coated with a synthetic cationic lipid can be introduced into cells by fusion.

The present invention further makes it possible to genetically engineer human hematopoietic progenitor cells in such a manner that they produce, *in vitro* or *in vivo* osteoclasts which produce polypeptides, hormones and proteins not normally produced in human osteoclasts in biologically significant amounts or produced in small amounts but in situations in which regulatory expression would lead to a therapeutic benefit. For example, the hematopoietic stem cells could be engineered with a gene that expresses a molecule that specifically inhibits bone resorption, but does not otherwise interfere with osteoclasts binding to bone. Alternatively the cells could be modified such that a protein normally expressed will be expressed at much lower levels. These products would then be secreted into the surrounding media or purified from the cells. The human osteoclast cells formed in this way can serve as continuous short term or long term production systems of the expressed substance.

This technology may be used to produce additional copies of essential genes to allow augmented expression by the osteoclasts of certain gene products *in vivo*. These genes can be, for example, hormones, matrix proteins, cell membrane proteins, cytokines, adhesion molecules, "rebuilding" proteins important in tissue repair. The expression of the exogenous genetic material *in vivo*, is often referred to as "gene therapy." Disease states and procedures for which such treatments have application include genetic disorders and diseases of bone and the immune system. Cell delivery of the transformed cells may be effected using various methods and includes infusion and direct depot injection into periosteal, bone marrow and subcutaneous sites.

In addition, as hereinabove described, the transduced cells may be used for *in vitro* production of desired protein(s). The transduced cells may further be used in screening assays for drug discovery.

In a preferred embodiment, hematopoietic stem cells are transduced with a retroviral vector containing a calcium response element, tartrate resistant acid phosphatase (TRAP) promoter and a cDNA encoding for anti-osteoclast factor.

After modification of the hematopoietic cells as described herein and induction of differentiation into osteoclasts, the mixture of hematopoietic, osteoclast and mesenchymal stem cells may be separated to obtain a population of cells largely consisting of the transduced osteoclasts. This may be accomplished
5 by positive and/or negative selection of transduced hematopoietic cells using antibodies to identify hematopoietic cell surface markers.

The above description of the invention and the following examples are by way of illustration only. Other permutations and practices of the invention will be
10 readily envisaged by one of ordinary skill in the art by view of the above in conjunction with the appended drawings. Therefore, such permutations and variations are within the scope of the present invention.

Examples

15

Statistical Analysis All data were analyzed by a paired t test. Samples were run in triplicates and data shown were means + SE.

Example 1

20

Cell isolation Bone marrow samples were collected from healthy human donors at the Johns Hopkins University under an Institutional Review Board-approved protocol. Mesenchymal stem cells (MSCs) were isolated and cultured according to known methods (Majumdar et al., 1998). When the cultures reached 90% confluence, adherent cells were recovered by the addition of a solution
25 containing 0.25% trypsin-EDTA (Life Technologies) and replated at a density of 1×10^6 cells per 185 cm² flask as passage-1 cells.

CD34+ cells, isolated from bone marrow of healthy patients were obtained from Poietic Technologies, Gaithersburg, MD). They were immunopurified to
30 95% purity using antibody to CD34 conjugated to magnetic beads (CD34+ cell separation column: Miltenyi Biotec, Auburn, CA Ab:QBEND.10) and cryopreserved.

Co-cultures MSCs of early passage (2 to 3) were plated at 3×10^3 cells per cm^2 and cultured until sub-confluency in DMEM LG+10% FBS. CD34⁺ cells were then added at 5×10^4 cells per cm^2 to MSC cultures. Cultures were maintained for 3 weeks at 37°C in an atmosphere of 95% air/5% CO₂ and fed every three days with the medium. Because most of the CD34⁺ cells remained non-adherent, half of the culture medium was gently aspirated without agitating non-adherent cells and replaced with fresh medium.

TRAP staining After three weeks, cells were fixed in 60% acetone in citrate buffer (pH 5.4) for 30 seconds, washed twice with distilled water, air-dried, and stained for tartrate-resistant acid phosphatase (TRAP), a widely-used cytochemical marker of mouse osteoclasts (Wijngaert and Burger, 1986), using a commercially available kit (Sigma, St. Louis, MO). Stained cultures were examined under light microscopy at a magnification of X200. TRAP-positive (red-staining) multinucleated (three or more nuclei) osteoclasts in each well were counted by manually scanning across the entire well in a systematic fashion (Mbalaviele et al., 1998).

CD34⁺ cells cultured in the absence of MSCs or cytokines/growth factors failed to form multinucleated cells expressing tartrate-resistant acid phosphatase (TRAP+MNCs) (Figure 1A). MSCs supported the differentiation of co-cultured CD34⁺ cells towards osteoclast cells (Figure 1B). In contrast, human skin fibroblasts (SK1087 cells) or human kidney cells (293T) failed to support the formation of TRAP+MNCs (Figures 1C and 1D, respectively). Treatment of CD34⁺ cells with IL-1, IL-3 and GM-CSF in the absence of MSCs yielded the formation of osteoclast cells, but the numbers of these cells were lower compared to those present in CD34⁺/MSCs co-cultures (data not shown).

Example 2

Immunocytochemistry CD34⁺ cells and MSCs co-cultured for three weeks were fixed with a fresh solution of 3.7% (wt./vol.) formaldehyde in PBS, blocked with 1.5% horse serum, incubated with the monoclonal antibody to human integrin $\alpha\text{v}\beta 3$ (Dr. Horton, The Rayne Institute, London, England) at 1:50 dilution in PBS

containing 0.15% horse serum. Cells were then incubated with glucose oxidase-conjugated goat secondary antibody using Vectastain ABC-GO kit (Vector Laboratories, Inc., Burlingame, CA). All incubations were performed at room temperature for 30 minutes followed by washing 3 times with PBS. Results are shown in Figure 2. Osteoclast cells (black arrow) and their precursors (arrowhead) were stained by vitronectin receptors antibody but not by the control antibody. MSCs did not show any staining with the receptor antibody. Multinucleated cells expressing low levels of vitronectin receptors were seen in co-cultures (Figure 2B, white arrow).

RNA Preparation and RT-PCR Total RNA was extracted from MSC cultures or co-cultures of CD34⁺ and MSCs using High Pure RNA Isolation kit purchased from Boehringer Mannheim (Indianapolis, IN). PCR was performed for 30 cycles either on reverse-transcribed single-strand cDNA from total RNA preparations (1 µg) using GeneAmp RT-PCR method under the conditions described by the supplier (Perkin Elmer, Foster City, CA) with the following modifications: denaturation at 95°C for 20 seconds, annealing at 55°C for 20 seconds, polymerization at 72°C for 30 seconds and elongation at 72°C for 10 minutes. The upstream and downstream primers, respectively, were designed as follows: TRAP: 5'-CGATCACAATCTGCAGTACC-3' (SEQ ID NO: 1) and 5'-ACCCAGTGAGTCTTCAGTCC-3' (SEQ ID NO:2), PCR product size=150 bp; calcitonin receptor (CTR): 5'-TTTCCAGGGCTTCTTTGTT-3' (SEQ ID NO:3) and 5'-CTTGGTTGTTGGCTGGTTC-3' (SEQ ID NO:4), PCR product size-205 bp. PCR products were size-fractionated by 1% agarose gel electrophoresis.

RT-PCR analysis showed expression of TRAP and calcitonin receptors by cells derived from CD34⁺ cells, the osteoclast lineage (Lanes 2 and 3 of Figure 3). These proteins were not expressed by MSCs.

Example 3

Pit formation assay To further confirm the presence of osteoclast cells in the cultures, the ability of cells produced in co-cultures of CD34⁺ cells and MSCs to form pits was examined, an indication of osteoclastic bone resorption. Elephant smoothed tusk slices were sterilized with 200-proof ethanol, air-dried and

washed several times with PBS. MSCs were plated at 3×10^3 cells per cm^2 on the tusk slices ($4 \times 4 \times 0.1$ mm) in 96-well plates and cultured for a week in hMSC medium pH 7.4. CD34^+ cells were then added at 5×10^4 cells per cm^2 . Artificial bone analog, osteologic discs (Millenium Biologix Inc., Ontario, Canada) were used to establish the optimal conditions of the assay. At the end of the 3-week culture period, the cells on slices were stained for TRAP, counted, then removed with 0.1 M NaOH in water followed by ultrasonication for 2 minutes. The bone slices were stained for 5 minutes with 1% toluidine blue prepared in distilled water containing 1% sodium borate. The numbers of pits were counted under light microscope as previously described (Prallet et al., 1992). Authentic resorption pits were found on bone analog as well as on elephant tusk slices when CD34^+ cells were co-cultured with MSCs (Figure 4).

Example 4

Effects of cell-cell interactions CD34^+ cells and MSCs were either cultured in the same compartment (contact co-cultures) or in two different compartments (non-contact co-cultures) separated by culture proof inserts with a $0.45 \mu\text{m}$ porous membrane (Becton Dickinson, Bedford, MA) to investigate physical cell-cell interactions. The results shown in Figure 5A indicated that osteoclast cells were formed in the contact cultures, and there were about two-thirds less osteoclast cells in non-contact cultures, regardless of in which compartment (top or bottom) the MSCs or CD34^+ cells were seeded.

Example 5

Interactions through secreted factors CD34^+ cells were co-cultured with MSCs. Conditioned media were collected after two days from MSCs cultured in the absence or presence of CD34^+ cells. The protein levels of cytokines IL- 1α , IL-6, IL-11, granulocyte/macrophage-colony stimulating factor (GM-CSF) and leukaemia inhibitory factor (LIF) secreted into the hematopoietic stem cell and mesenchymal stem cell co-cultures were measured using a Quantikine kit (R&D Systems, Minneapolis, MN).

Figure 5B shows the levels of IL-6, IL-11 or LIF in co-cultures of CD34⁺ cells and MSCs. The levels of IL-1 α and GM-CSF were undetectable. The levels of IL-6 and LIF were about 10 times greater in co-cultures of MSCs and CD34⁺ cells compared to MSCs cultures without CD34⁺ cells. The levels of IL-1 α and GM-CSF were undetectable.

Example 6

Effects of osteogenic MSCs (differentiated MSCs) Osteogenic differentiation of MSCs was induced by treating MSCs with osteogenic supplement (OS) containing 100mM dexamethasone 10mM β -glycerophosphate and 50 μ M L-ascorbic acid-2 phosphate for 2, 3, 4, 5, 10 or 13 days. At the time of the co-culture with CD34⁺ cells, OS containing medium was replaced with medium without OS. Figures 6A and 6B show that osteogenic MSCs inhibited TRAP⁺MNC formation compared to MSCs. Figure 6C shows the levels of IL-6, IL-11 or LIF in co-culture. The levels of IL-6, IL-11 and LIF were higher in co-cultures of MSCs with CD34⁺ cells compared to cocultures of osteogenic MSCs and CD34⁺ cells.

RNA preparation and RT-PCR Total RNA isolation and RT-PCR analysis were carried out as described in example 2. The upstream and downstream primers, respectively, for OPG were 5'-ACCACTACTACACAGACAGC-3' (SEQ ID NO:5) and 5'-AGGAGACCAAAGACACTGCA-3' (SEQ ID NO:6) ; for OPGL 5'-TTCTATTTTCAGAGCGCAGAT-3' (SEQ ID NO:7) and 5'-AGTCATGTTGGAGATCTTGG-3' (SEQ ID NO:8). Figure 6D shows that treatment of MSCs with OS for 4, 8, or 15 days decreased the expression of OPGL mRNA. In contrast, treatment of MSCs with OS for 4, 8 or 15 days increased the expression of OPG mRNA.

Example 7

Transduction of CD34⁺ cells The construction of the retroviral vector MGIN expressing enhanced green fluorescent protein (EGFP) gene and the production of the retroviral supernatants has been previously reported (Cheng et al., 1997). Briefly, MGIN is a murine embryonic stem cell virus-based vector

containing EGFP gene and the internal ribosome entry site (IRES). Amphotropic supernatants produced by PA317 packaging cells were made from selected producers after infection by BOSC23 ecotropic viral stocks. For transduction, previously frozen vector supernatants were mixed at a 1:1 ratio with medium containing CD34⁺ cells in the presence of 8 µg/ml polybrene (Sigma, St. Louis, MO), interleukin-3 (IL-3) and IL-6 (10 ng/ml, each), stem cell factor (SCF) and Flk2 (FL) (100 ng/ml, each). Control cells were transduced with non-EGFP-expressing vector. The transduction mixture was then centrifuged at 1800 g at 32-35°C. After a 4-hour "spinoculation," cells were washed once and cultured for 24 hours in medium containing the above growth factors. After the transduction was repeated once more, a cell fraction was analyzed by flow cytometry to test transduction efficiency while the remaining cells were used for the osteoclastogenesis assay. The transduced CD34⁺ cells were co-cultured for three weeks without added factors. The co-cultures were stained for TRAP as described in example 1. Cells were visualized with a fluorescence microscope to detect EGFP or under a light microscope to detect TRAP.

Flow cytometry analysis results are shown in Figure 7. FACS analysis showed that 30% of CD34⁺ cells transduced with EGFP vector expressed the transgene (Figure 7 B). Co-cultures of transduced CD34⁺ cells with MSCs yielded osteoclasts expressing both EGFP (Figure 7C) and TRAP (Figure 7D).

Example 8

Detection of vitronectin receptor by ELISA MSCs were plated at 3×10^3 cells per cm² and cultured in hMSC medium until subconfluency. CD34⁺ cells were then added at 2.5×10^4 cells per cm² to MSC cultures. Cultures were maintained for 1, 2 or 3 weeks. Cells were fixed with 3.6% formaldehyde and blocked with 10% goat serum for 30 minutes, followed by incubation with the antibody to the vitronectin receptor ($\alpha v \beta 3$). With extensive washes between additions, a biotin-labeled secondary antibody was added (goat anti-mouse IgG; Pierce, Rockford, IL) followed by streptavidin-conjugated β -galactosidase (Gibco BRL) for 1 hour. The soluble substrate chlorophenol red- β -D-galactopyranoside (CPRG; Boehringer Mannheim, Indianapolis, IN) was then added, and color

development was read at 570 nm after 30 minutes. Control primary antibodies were also run for each treatment and this background measurement was subtracted from the readings for each well. ELISA results are presented in Figure 8 as mean absorbance units, with background subtracted. The vitronectin receptor ($\alpha v\beta 3$) was detected by the specific antibody but not by the control antibody (Figure 8A). $\alpha v\beta 3$ expression by the osteoclast cell lineage was induced after 1 week of co-cultures and was less expressed by MSCs (Figure 8B).

Example 9

Hematopoietic stem cells as therapeutic targets Hematopoietic stem cells (HSCs) are transduced with a retroviral vector containing a calcium response element, tartrate resistant acid phosphatase (TRAP) promoter and a cDNA encoding for anti-osteoclast factor (antisense, inhibitor, apoptotic factor) (see Figure 8). TRAP promoter has been used to specifically target the osteoclast cell lineage. Transduced-HSCs are infused to osteoporotic patients. They home to bone marrow, engraft and differentiate into osteoclasts. The osteoclasts respond to local high calcium concentrations generated by excessive bone resorption (steps 1 and 2), by expressing an anti-osteoclast factor (step 3) which stops osteoclast activity (step 4). One such factor is Osteoprotegerin (OPG) which when secreted has been demonstrated to decrease bone resorption (osteopetrosis) and appears to generally block osteoclastogenesis (Simonet (1997)). The transduced hematopoietic cells can also be induced to differentiate into osteoclasts ex vivo, and then infused to osteoporotic patients.

Calcium-response element (CaRE) was cloned under the control of CMV promoter within pGL3 vector that contains luciferase as a reporter gene (Promega Madison WI) to yield pGL3-CaRE. pGL3 or pGL3-CaRE were transfected into human kidney cell line 293T using a calcium phosphate co-precipitation method. Cells were then cultured in the presence of increasing concentrations of calcium. Luciferase activity was assessed using Dual-Luciferase™ reporter assay system (Promega). The data show that calcium as low as 0.1mM induced luciferase activity in pGL3-CaRE but not in pGL3 (Figure 9B).

Example 10

Effect of serum in culture medium Co-cultures of MSCs and CD34+ cells were studied to determine the effect of serum on the production of osteoclasts. Co-cultures of CD34+ cells and MSCs were cultured, under the conditions described above, however the media tested were DMEM-low glucose with 10% FBS and DMEM-high glucose without serum. Control cultures contained human kidney cell line 293 cells co-cultured with CD34+ cells in DMEM-high glucose with and without serum. The cultures were terminated at 12 days rather than 21 days as in the experiments in the foregoing examples.

10

The results shown in Figure 10 demonstrate that co-culturing MSCs and CD34+ cells in DMEM-high glucose without FBS yielded a greater number of osteoclasts than co-culture in the standard DMEM-LG with 10% FBS.

CITED LITERATURE

Caplan AI, Bruder SP: Cell and molecular engineering of bone regeneration, in Lanza R, Langer R, Chick W (eds): Textbook of Tissue Engineering, Georgetown, Landes Company, 1997, p 603

5

Cheng L, Du C, Murray D, Tong X, Zhang YA, Chen BP, Hawley RG: A GFP reporter system to assess gene transfer and expression in human hematopoietic progenitor cells. *Gene Therapy* 4: 1013 (1997)

- 10 James IE, Dodds RA, Lee-Rykaczewski E, Eichman CF, Connor JR, Hart T, Maleef BE, Lackman RD, Gowen M: Purification and characterization of fully functional human osteoclast precursors. *J. Bone Min. Res.* 11: 1608 (1996)

- 15 Kong YY, Yoshida H, Sarosi I, Tan HL, Timms E, Capparelli C, Morony S, Oliveira-dos-Santos AJ, Van G, Itie A, Khoo W, Wakeham A, Dunstan CR, Lacey DL, Mak TW, Boyle WJ, Penninger JM: OPGL is a key regulator of osteoclastogenesis, lymphocyte development and lymph-node organogenesis. *Nature* 397(6717):315-23 (1999)

- 20 Lacey DL, Timms E, Tan HL, Kelley MJ, Dunstan CR, Burgess T, Elliott R, Colombero A, Elliott G, Scully S, Hsu H, Sullivan J, Hawkins N, Davy E, Capparelli C, Eli A, Qian YX, Kaufman S, Sarosi I, Shalhoub V, Senaldi G, Guo J, Delaney J, Boyle WJ: Osteoprotegerin ligand is a cytokine that regulates osteoclast differentiation and activation. *Cell* 93(2):165-76 (1998)

25

McLachlin et al. Progress in Nucleic Acid Research and Molecular Biology, 38:91-135(1990)

- 30 Majumdar MK, Thiede MA, Mosca JD, Moorman M, Gerson S: Phenotypic and functional comparison of cultures of marrow-derived mesenchymal stem cells (MSCs) and stromal cells. *J Cell. Physiol.* 176(1):57-66 (1998)

Matayoshi A, Brown C, DiPersio JF, Haug J, Abu-Amer Y, Liapis H, Kuestner R, Pacifici R: Human blood-mobilized hematopoietic precursors differentiate into

osteoclasts in the absence of stromal cells. *Proc. Natl. Acad. Sci. USA* 93, 10785 (1996)

5 Mbalaviele G, Orcel P, Morieux C, Nijweide PJ, Devemejoul MC: Osteoclast formation from human cord blood mononuclear cells co-cultured with mice embryonic metatarsals in the presence of M-CSF. *Bone* 16: 171 (1995)

10 Mbalaviele G, Nishimura R, Myoi A, Niewolna M, Reddy SV, Chen D, Feng J, Roodman D, Mundy GR, Yoneda T: Cadherin-6 mediates the heterotypic interactions between the hemopoietic osteoclast cell lineage and stromal cells in a murine model of osteoclast differentiation. *J. Cell Biol* 141(6):1467-1476 (1998).

15 Mundy GR, Boyce BF, Yoneda T, Bonewald LF, Roodman GD: Cytokines and bone remodeling, in Marcus R, Feldman D, Kelsey J (eds): Osteoporosis, San Diego, Academic Press, 1995, p 302

20 Prallet B, Male P, Neff L, Baron R: Identification of a functional mononuclear precursor of the osteoclast in chicken medullary bone marrow cultures. *J Bone Min. Res.* 7: 405 (1992)

Sarma U, Flanagan AM: Macrophage colony-stimulating factor induces substantial osteoclast generation and bone resorption in human bone marrow cultures. *Blood*, 88: 2531 (1996)

25 Quinn JMW, Fujikawa Y, McGee JO'D, Athanasou NA: Rodent osteoblast-like cells support osteoclastic differentiation of human cord blood monocytes in the presence of M-CSF and 1,25 Dihydroxyvitamin D3. *Int. J. Biochem. Cell Biol.* 29: 173 (1997)

30 Simonet et al. Osteoprotegerin: A Novel Secreted Protein Involved in the Regulation of Bone Density. *J. Biol. Chem.* 268(1):377-384 (1993)

Takahashi S, Reddy SV, Dallas M, Devlin R, Chou JY, Roodman GD: Development and characterization of a human marrow stromal cell line that enhances osteoclast-like cell formation. *Endocrinol.* 136: 1441 (1995)

- 5 Takahashi S. et al. *Endocrinology* 123:2600-2602 (1988)

Teitelbaum SL, Tondravi MM, Ross FP: Osteoclast biology, in Marcus R, Feldman D, Kelsey J (eds): Osteoporosis, San Diego, Academic Press, 1995, p 61

- 10 Wijngaert FB, Burger EH: Demonstration of tartrate-resistant acid phosphatase in undecalcified, glycomethacrylate-embedded mouse bone: a possible marker for (pre-osteoclast identification *J Histo. Cytol.* 34:1317 (1986)

CLAIMS

1. A method for producing osteoclasts *in vitro* comprising co-culturing
5 mesenchymal stem cells with hematopoietic stem cells such that the hematopoietic
stem cells produce osteoclasts.
2. The method of claim 1 wherein co-culturing is done in the absence of
exogenous cytokines.
- 10 3. The method of claim 1 wherein the hematopoietic stem cells are CD34+
cells.
4. The method of claim 1 wherein the hematopoietic stem cells and the
15 mesenchymal stem cells are in contact.
5. A method of producing genetically modified osteoclasts, comprising
transducing hematopoietic progenitor cells with exogenous genetic material; and
culturing the transduced hematopoietic cells in the presence of mesenchymal stem
20 cells to induce differentiation of the transduced hematopoietic cells into osteoclasts
which contain the exogenous genetic material.
6. The method of claim 5 wherein the transduced hematopoietic stem cells
are induced to differentiate *in vitro*.
- 25 7. The method of claim 5 wherein the transduced hematopoietic stem cells
are induced to differentiate *in vivo*.
8. A method of obtaining genetically modified osteoclasts comprising
30 transducing hematopoietic progenitor cells with exogenous genetic material and
placing the transduced hematopoietic cells under conditions suitable for
differentiation of the hematopoietic stem cells into osteoclast cells which contain
the exogenous genetic material.

9. The method of claim 8 wherein the hematopoietic cells are differentiated *in vivo*.

10. The method of claim 8 wherein the hematopoietic cells are administered *in vivo* with human mesenchymal stem cells.

11. A composition comprising genetically modified osteoclasts.

12. The composition of claim 11 wherein the genetically modified osteoclasts are from a non-transformed cell line.

13. A composition comprising genetically modified osteoclast precursor cells.

14. The composition of claim 13 wherein the genetically modified osteoclast precursor cells are from a non-transformed cell line.

15. A method for decreasing differentiation of hematopoietic stem cells into osteoclasts comprising co-culturing differentiating mesenchymal stem cells with hematopoietic stem cells such that osteoclastogenesis is reduced.




Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 6




Figure 7




Figure 8




Figure 9




Figure 10

SEQUENCE LISTING

<110> Osiris Therapeutics, Inc.

<120> Regulation of Hematopoietic Stem Cell Differentiation
by the Use of Human Mesenchymal Stem Cells

<130> 640100-333

<140>

<141>

<150> 60/099,233

<151> 1998-09-04

<150> 60/088,431

<151> 1998-06-08

<160> 8

<170> PatentIn Ver. 2.0

<210> 1

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: PCR Primer

<400> 1

cgatcacaat ctgcagtacc

20

<210> 2

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: PCR Primer

<400> 2

acccagtgag tcttcagtcc

20

<210> 3

<211> 19

<212> DNA

<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: PCR Primer

<400> 3
tttccagggc ttctttgtt 19

<210> 4
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: PCR Primer

<400> 4
cttggttggtt ggctggttc 19

<210> 5
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: PCR Primer

<400> 5
accactacta cacagacagc 20

<210> 6
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: PCR Primer

<400> 6
aggagaccaa agacactgca 20

<210> 7
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: PCR Primer

<400> 7

ttctattttca gagcgagat

20

<210> 8

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: PCR Primer

<400> 8

agtcacgttg gagatcttgg

20



PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 5/06, 5/08, 5/10, A61K 35/12, 48/00		A3	(11) International Publication Number: WO 99/64565
			(43) International Publication Date: 16 December 1999 (16.12.99)
(21) International Application Number: PCT/US99/12847 (22) International Filing Date: 8 June 1999 (08.06.99) (30) Priority Data: 60/088,431 8 June 1998 (08.06.98) US 60/099,233 4 September 1998 (04.09.98) US (71) Applicant: OSIRIS THERAPEUTICS, INC. [US/US]; 2001 Aliceanna Street, Baltimore, MD 21231-2001 (US). (72) Inventors: THIEDE, Mark, A.; 319 Pond View Court, Forest Hill, MD 21050 (US). MBALAVIELE, Gabriel; Apartment 1G, 8488 Greystone Lane, Columbia, MD 21045 (US). (74) Agents: OLSTEIN, Elliot, M. et al.; Carella, Byrne, Bain, Gilfillan, Cecchi, Stewart & Olstein, 6 Becker Farm Road, Roseland, NJ 07068 (US).			(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> (88) Date of publication of the international search report: 24 February 2000 (24.02.00)
(54) Title: REGULATION OF HEMATOPOIETIC STEM CELL DIFFERENTIATION BY THE USE OF HUMAN MESENCHYMAL STEM CELLS			
(57) Abstract The invention relates to the induction of hematopoietic stem cells to differentiate into osteoclasts by culturing hematopoietic stem cells with human mesenchymal stem cells, and, in a preferred embodiment, using no exogenous cytokines. Differentiation of the mesenchymal stem cells into osteoblasts inhibited the differentiation of hematopoietic stem cells into osteoclasts. In addition, hematopoietic stem cells can be genetically engineered to carry genes of interest particularly for the expression of physiologically active proteins. In the presence of mesenchymal stem cells, the transduced cells carry the new genetic material and express gene products that can be used to modulate bone resorption.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

INTERNATIONAL SEARCH REPORT

Inter. Application No
PCT/US99/12847

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N5/06 C12N5/08 C12N5/10 A61K35/12 A61K48/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 670 351 A (EMERSON STEPHEN G ET AL) 23 September 1997 (1997-09-23) column 10, line 51 -column 11, line 67 claims 1,18,50,108 ---	8,11-14
A	WO 95 02040 A (GSF FORSCHUNGSZENTRUM UMWELT ;THALMEIER KARIN (DE); DOERMER PETER) 19 January 1995 (1995-01-19) the whole document ---	1-15
A	WO 96 32469 A (RODAN SEVGI B ;MERCK & CO INC (US); RODAN GIDEON A (US); WESOLOWSK) 17 October 1996 (1996-10-17) the whole document ---	1-15
A	US 5 486 359 A (CAPLAN ARNOLD I ET AL) 23 January 1996 (1996-01-23) the whole document ---	1-15
-/--		

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

14 December 1999

Date of mailing of the international search report

20/12/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Fernandez y Branas, F

INTERNATIONAL SEARCH REPORT

International Application No

PC 99/12847

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>TAKAHASHI S ET AL: "Development and characterization of a human marrow stromal cell line that enhances osteoclast -like cell formation." ENDOCRINOLOGY, (1995 APR) 136 (4) 1441-9. , XP000857459 cited in the application page 1443, column 1, paragraph 3 page 1445, column 2, paragraphs 1,2 -----</p>	1-15
T	<p>MBALAVIELE G ET AL: "Human mesenchymal stem cells promote human osteoclast differentiation from CD34+ bone marrow hematopoietic progenitors." ENDOCRINOLOGY, (1999 AUG) 140 (8) 3736-43. , XP000857472 the whole document -----</p>	1-15

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 99/ 12847

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 5 (partially), 7, 8 (partially), 9-10, 15 (partially) are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/12847

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 5670351	A	23-09-1997	US 5399493 A	21-03-1995
			AU 687674 B	26-02-1998
			AU 5059296 A	11-07-1996
			AU 665525 B	11-01-1996
			AU 9175091 A	22-07-1992
			CA 2100268 A	18-06-1992
			EP 0575350 A	29-12-1993
			JP 11221074 A	17-08-1999
			JP 6505151 T	16-06-1994
			WO 9211355 A	09-07-1992
			US 5635386 A	03-06-1997
			US 5437994 A	01-08-1995
			US 5605822 A	25-02-1997
			US 5670147 A	23-09-1997
			US 5646043 A	08-07-1995
			AT 148502 T	15-02-1997
			CA 2062741 A	16-12-1990
			DE 69029856 D	13-03-1997
			DE 69029856 T	04-09-1997
			DK 477290 T	21-07-1997
			EP 0477290 A	01-04-1992
			EP 0753574 A	15-01-1997
			ES 2097149 T	01-04-1997
			JP 4506153 T	29-10-1992
			WO 9015877 A	27-12-1990
			US 5459069 A	17-10-1995
			US 5763266 A	09-06-1998
			US 5888807 A	30-03-1999
WO 9502040	A	19-01-1995	DE 4322570 C	16-06-1994
			AU 686218 B	05-02-1998
			AU 7491494 A	06-02-1995
			CA 2166707 A,C	19-01-1995
			CN 1126489 A	10-07-1996
			CZ 9600056 A	17-07-1996
			WO 9502039 A	19-01-1995
			EP 0707634 A	24-04-1996
			FI 960064 A	05-01-1996
			HU 73380 A	29-07-1996
			JP 8508413 T	10-09-1996
			NO 960050 A	05-01-1996
			US 5658761 A	19-08-1997
WO 9632469	A	17-10-1996	US 5719058 A	17-02-1998
			CA 2217816 A	17-10-1996
			EP 0821727 A	04-02-1998
			US 5856186 A	05-01-1999
US 5486359	A	23-01-1996	US 5733542 A	31-03-1998
			US 5811094 A	22-09-1998
			US 5837539 A	17-11-1998
			AU 669850 B	27-06-1996
			AU 2252492 A	12-01-1993
			CA 2111845 A	23-12-1992
			EP 0592521 A	20-04-1994
			JP 7500001 T	05-01-1995
			WO 9222584 A	23-12-1992



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : C12N 5/06, 5/08, 5/10, A61K 35/12, 48/00		A3	(11) International Publication Number: WO 99/64565
		(43) International Publication Date: 16 December 1999 (16.12.99)	
(21) International Application Number: PCT/US99/12847 (22) International Filing Date: 8 June 1999 (08.06.99) (30) Priority Data: 60/088,431 8 June 1998 (08.06.98) US 60/099,233 4 September 1998 (04.09.98) US (71) Applicant: OSIRIS THERAPEUTICS, INC. [US/US]; 2001 Aliceanna Street, Baltimore, MD 21231-2001 (US). (72) Inventors: THIEDE, Mark, A.; 319 Pond View Court, Forest Hill, MD 21050 (US). MBALAVIELE, Gabriel; Apartment 1G, 8488 Greystone Lane, Columbia, MD 21045 (US). (74) Agents: OLSTEIN, Elliot, M. et al.; Carella, Byrne, Bain, Gilfillan, Cecchi, Stewart & Olstein, 6 Becker Farm Road, Roseland, NJ 07068 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> (88) Date of publication of the international search report: 24 February 2000 (24.02.00)	
(54) Title: REGULATION OF HEMATOPOIETIC STEM CELL DIFFERENTIATION BY THE USE OF HUMAN MESENCHYMAL STEM CELLS			
(57) Abstract The invention relates to the induction of hematopoietic stem cells to differentiate into osteoclasts by culturing hematopoietic stem cells with human mesenchymal stem cells, and, in a preferred embodiment, using no exogenous cytokines. Differentiation of the mesenchymal stem cells into osteoblasts inhibited the differentiation of hematopoietic stem cells into osteoclasts. In addition, hematopoietic stem cells can be genetically engineered to carry genes of interest particularly for the expression of physiologically active proteins. In the presence of mesenchymal stem cells, the transduced cells carry the new genetic material and express gene products that can be used to modulate bone resorption.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

5

10 **REGULATION OF HEMATOPOIETIC STEM CELL DIFFERENTIATION
BY THE USE OF HUMAN MESENCHYMAL STEM CELLS**

 The present invention relates to hematopoietic stem cells and more particularly to a process and composition for differentiating human hematopoietic stem cells into a committed osteoclast lineage in the absence of exogenous
15 osteoclastogenesis-inducing factors. The present invention further relates to genetically modified hematopoietic stem cells in the presence of human mesenchymal stem cells, such that when the hematopoietic stem cells differentiate into osteoclasts, the osteoclasts are able to express the product of the transduced
20 gene.

Background of the Invention

 Osteoclasts are terminally differentiated hematopoietic cells responsible for physiological bone resorption. Dysfunction in osteoclast differentiation and/or
25 activity is the cause of a variety of human metabolic bone diseases including osteoporosis (for review see Teitelbaum et al., 1995). Osteoporosis, or progressive bone loss, is attributed to a shift in the critical balance of osteoclast and osteoblast activities responsible for proper skeletal integrity. Osteoblasts, which build bone, are one terminally differentiated lineage descending from mesenchymal stem cells.

30

 Hematopoietic progenitors in bone marrow are driven into the osteoclast cell lineage under the influence of local regulatory factors. Diverse cell types including stromal cells, which arise from mesenchymal stem cells (MSCs), are present in bone marrow. Stromal cells have been shown to produce factors that
35 regulate osteoclast physiology (for review see Mundy et al., 1995). Histological

pictures of bone show close vicinity between marrow stromal cells and hematopoietic cells. Further, the relationship between the differentiated status of bone marrow stromal cells and their potential to regulate neighboring osteoclast progenitors is poorly documented. The importance of heterotypic cell-to-cell interactions mediated by cadherin-6 between stromal cells and osteoclast progenitors using a mouse model of osteoclastogenesis has been reported (Mbalaviele et al., 1998). Several studies demonstrated that osteoclast differentiation required the presence of stromal/osteoblast cells (Takahashi et al., 1988; Mbalaviele et al., 1995). Recent studies using cells derived from rodent marrow have demonstrated that osteoprotegerin ligand (OPGL) expressed on the membrane of stromal cells is a key factor of osteoclastogenesis (Lacey et al. 1998; Kong et al. 1999). However, it has also been reported that human osteoclast cells differentiated from progenitor cells in the presence of cocktails of cytokines and that stromal cells were not necessary (Matayoshi et al., 1996). The formation of human osteoclast cells (Ocl) displaying bone-resorbing activity has been reported (Takahashi et al., 1995; James et al., 1996; Sarma et al., 1996; Quinn et al., 1997), however, critical events which take place during osteoclastogenesis are unclear.

Bone is a composite matrix consisting of both inorganic and organic elements. The organic phase contains proteins, while the inorganic component contains calcium salts. It is known that during osteoclastic bone resorption, dissolution of the inorganic phase precedes that of the organic phase and that high concentrations of calcium (>40mM) are released locally from bone matrix. In turn, increased calcium concentrations have a negative feedback on osteoclast function, a regulatory mechanism to prevent excessive bone resorption. Calcium is known to regulate the expression of several genes. Recently, a calcium response element that enhances human keratin-1 gene expression after exposure of keratinocytes to extracellular calcium (0.6 mM) has been characterized.

Osteoclasts are not easily isolated due to their low cell number *in vivo*, their fragility and tendency to adhere to other bone cells. Thus, there is a need for a method of efficiently producing quantities of osteoclasts to facilitate, for example, the design of effective therapeutics aimed at preventing abnormal osteolysis.

Accordingly, it is an object of the present invention to provide an efficient method of producing osteoclasts *in vitro*.

It is a further object of the invention to provide a method to produce and
5 maintain osteoclasts *in vitro* without the addition of exogenous growth factors in numbers sufficient to provide an ongoing source of osteoclasts, for example, for administering osteoclasts to an individual in need thereof.

Accordingly, it is an object of the present invention to regulate
10 hematopoietic stem cell differentiation towards osteoclast cells in the absence of exogenous growth factors, cytokines and hormones.

It is a still further object of the present invention to provide a method of producing osteoclasts from genetically modified hematopoietic cells such that
15 when the hematopoietic cells differentiate into osteoclasts, the osteoclasts express the gene product of interest.

Summary of the Invention

The present inventors have discovered that mesenchymal stem cells support
20 osteoclast cell differentiation of hematopoietic stem cells.

Thus in one aspect of the present invention, there is provided a method for inducing the differentiation of hematopoietic progenitor cells *in vitro* into osteoclast cells, comprising co-culturing the hematopoietic cells with human
25 mesenchymal stem cells.

When hematopoietic progenitor cells were co-cultured with mesenchymal stem cells, the hematopoietic progenitor cells differentiated towards osteoclast cells which were identified by the expression of markers characteristic of osteoclasts
30 including multinuclearity, calcitonin and vitronectin receptors, and more importantly, bone resorbing activity.

It has further been discovered that the differentiation of the hematopoietic stem cells occurred in the absence of exogenous (added) growth factors, cytokines

and hormones known to be required for inducing differentiation of hematopoietic stem cells.

Accordingly, one aspect of the present invention provides a method of
5 producing osteoclasts *in vitro* comprising co-culturing hematopoietic stem cells with human mesenchymal stem cells. In a preferred embodiment, the cells are co-cultured in the same culture vessel such that physical cell-to-cell interactions occur between the hematopoietic progenitor cells and the mesenchymal stem cells.

10 It has also been discovered that when hematopoietic stem cells, which have been modified to carry exogenous genetic material of interest, are co-cultured with mesenchymal stem cells, the transduced hematopoietic stem cells differentiated into osteoclasts that also carried the new genetic material. These transduced
15 osteoclast cells are able to express the exogenous gene product. Transduced osteoclast progenitors and the osteoclasts differentiated therefrom can be used for applications where treatment using such modified osteoclasts is beneficial, for example, in the alleviation of the effects of osteoporosis.

Accordingly, the present invention provides a method of obtaining
20 genetically modified osteoclasts, comprising transducing hematopoietic progenitor cells with exogenous genetic material and placing the transduced hematopoietic cells under conditions suitable for differentiation of the hematopoietic stem cells into osteoclast cells which contain the exogenous genetic material.

25 In one embodiment, the method of producing osteoclasts comprises co-culturing transduced hematopoietic stem cells with mesenchymal stem cells such that after differentiation of the hematopoietic stem cells into osteoclasts, the osteoclasts also contain the exogenous genetic material.

30 The invention further relates to hematopoietic stem cells that are transduced with a polynucleotide that encodes a product that downregulates osteoclast differentiation and/or activity, and the use thereof.

In a particularly preferred embodiment, the expressed gene product downregulates osteoclast differentiation and/or activity. Hematopoietic stem cells are transduced with a retroviral vector containing a tandem of a calcium response element, tartrate resistant acid phosphatase (TRAP) promoter (specifically
5 expressed in osteoclasts) and a nucleic acid sequence encoding an anti-osteoclast activity factor (for example, an antisense sequence to M-CSF, IL1 or IL6; an inhibitor such as an intracellular antibody to IL6, to M-CSF, to GM-CSF, to IL1, or to Leukemia Inhibitory Factor; or, an apoptotic factor). Expression of this sequence is triggered by the local increase in calcium concentration due to
10 osteoclastic bone resorption.

It has further been found that while mesenchymal stem cells positively regulated the differentiation of CD34⁺ progenitor cells towards osteoclast cells, MSCs that have begun differentiation into osteoblastic cells inhibited this process.
15 In the presence of osteogenic mesenchymal stem cells, the levels of OPGL expression decreased and expression of osteoprotegerin (OPG) were found to increase. Thus, it is also contemplated that culturing hematopoietic stem cells in the presence of differentiated mesenchymal stem cells may serve to downregulate osteoclast differentiation and/or activity. Accordingly, the co-cultured cell
20 populations of the invention may be used as part of a treatment to alleviate the symptoms of osteoporosis.

Brief Description of the Drawings

Figure 1A shows TRAP staining of CD34⁺ cells cultured in the absence of
25 mesenchymal stem cells (MSCs).

Figure 1B shows TRAP staining of CD34⁺ cells cultured in the presence of MSCs. Many multinucleated TRAP⁺ cells (TRAP+MNCs) (arrow) as well as mononuclear TRAP⁺ cells were formed in the presence of MSCs which appeared as a confluent layer of spindle-shaped cells (arrowhead). Small arrows and star
30 indicate clusters of nuclei.

Figure 1C shows staining of CD34⁺ cells cultured in the presence of human skin fibroblasts (SK1087 cells; arrowhead). No TRAP+MNCs were formed.

Figure 1D shows staining of CD34⁺ cells cultured in the presence of human kidney cells (293T; arrowhead). TRAP+MNCs were not formed.

Figure 2A shows immunocytochemical staining of co-cultures of CD34⁺ and MSCs with negative-control antibody.

Figure 2B shows immunocytochemical staining of co-cultures of CD34⁺ and MSCs with antibody recognizing vitronectin receptor.

Figure 3 shows RT-PCR of RNA purified from co-cultures of MSCs and CD34⁺ cells (lanes 2-3) or cultures of MSCs without CD34⁺ cells (lanes 4-5). Calcitonin receptors (lanes 2 and 4) and TRAP (lanes 3 and 5) were only detected in samples from co-cultures of MSCs and CD34⁺ cells. The conserved and unspliced region of the calcitonin receptor family was chosen to design primers. Lane 1, 1 kb DNA ladder; bottom, β 2-microglobulin.

Figure 4A shows pit formation in co-cultures of CD34⁺ cells and MSCs (arrows).

Figure 4B shows pit formation in cultures of CD34⁺ cells without MSCs.

Slices were stained with toluidine blue and visualized under light microscopy. X200 magnification.

Figure 4C shows quantitation of pits in co-cultures of CD34⁺ cells and MSCs (closed bar) or in cultures of CD34⁺ cells without MSCs (open bar). * $p < 0.01$ versus cultures of CD34⁺ cells without MSCs.

Figure 5A shows the effects of cell-to-cell interactions in osteoclast cell formation. CD34⁺ cells and MSCs were co-cultured in the same compartments (contact co-cultures, closed bar) or separated by means of 0.45 μ m filters (non-contact co-cultures): MSCs on top and CD34⁺ cells on the bottom, open bar; MSCs on the bottom and CD34⁺ cells on top, dashed bar). * $p < 0.01$ versus non-contact co-cultures.

Figure 5B shows quantitation of secreted factors (IL-6, IL-11, LIF) in co-cultures of CD34⁺ cells with MSCs. The levels of IL-6 and LIF but not IL-11 were about 10 times higher in co-cultures of MSCs and CD34⁺ cells compared to MSC cultures without CD34⁺ cells

Figure 6A and 6B show inhibition of TRAP+MNCs by osteogenic MSCs. Osteogenic differentiation of MSCs was induced by treatment with osteogenic supplement (OS) for 2, 3, 4, 5, 10 or 13 days prior to adding CD34+ cells. *p<0.01 versus co-cultures of OS-treated MSCs and CD34+ cells.

5 Figure 6C shows that co-cultures of MSCs and CD34+ cells produced higher levels of IL-6, IL-11 and LIF than co-cultures of osteogenic MSCs and CD34+ cells.

10 Figure 6D shows that osteogenic differentiation of MSCs is associated with a decrease in OPGL mRNA expression (left panel) and an increase in OPG mRNA expression (right panel).

Figure 7A shows flow cytometry analysis of CD34⁺ cells transduced with control vector.

15 Figure 7B shows flow cytometry analysis of CD34⁺ cells transduced with EGFP expression vector. Approximately 30% of CD34⁺ cells expressed EGFP.

Figure 7C shows co-cultures of MSCs with CD34⁺ cells transduced with EGFP expression vector. EGFP was expressed by osteoclasts as well as by mononuclear cells.

20 Figure 7D shows co-cultures of MSCs with CD34⁺ cells transduced with EGFP expression vector stained for TRAP. Fig. 7D is the light view of Fig. 7C.

Figure 8A shows the expression of vitronectin receptor ($\alpha_v\beta_3$) detected by ELISA. $\alpha_v\beta_3$ antibody, wells on left and closed bar; control IgG, wells on right and open bar.

25 Figure 8B shows the time-course expression of $\alpha_v\beta_3$. Co-cultures of CD34+ cells and MSCs (closed bars) or MSCs without CD34+ cells (open bars).

Figure 9A shows a strategy for direct regulation of osteoclast activity.

30 Figure 9B shows that the addition of calcium to cultured cells transfected with a calcium-inducible expression system induced expression of the reporter gene.

Figure 10 shows the effect of serum in culture medium on production of osteoclasts in co-cultures of MSCs and CD34+ cells.

Figure 10A shows osteoclast numbers in co-cultures of CD34+ cells and MSCs in DMEM-low glucose with 10% FBS, open bar; and in DMEM-high glucose without serum, solid bar.

Figure 10B show co-cultures of 293 cells and CD34+ cells in DMEM-high glucose with 10% FBS, solid bar; and in DMEM-high glucose without serum, dashed bar.

Detailed Description of the Invention

The present invention relates generally to the use of human mesenchymal stem cells to induce the differentiation of human CD34+ cells into osteoclast cells and compositions comprising human CD34+ cells and human mesenchymal stem cells. More particularly, the inventors found that human mesenchymal stem cells cultured in association with CD34+ cells are useful for inducing the differentiation of CD34+ cells into osteoclasts. Thus the CD34+ cells can be maintained and utilized as a source of osteoclasts.

In order to obtain subject human mesenchymal stem cells for the methods described herein, mesenchymal stem cells can be recovered from bone marrow or other mesenchymal stem cell source. Bone marrow cells may be obtained from iliac crest, femora, tibiae, spine, rib or other medullary spaces. Other sources of human mesenchymal stem cells include embryonic yolk sac, placenta, umbilical cord, fetal and adolescent skin, and blood. The presence of mesenchymal stem cells in the culture colonies may be verified by specific cell surface markers which are identified with unique monoclonal antibodies, see, e.g., U.S. Patent No. 5,486,359. These isolated mesenchymal cell population display epitopic characteristics associated only with mesenchymal stem cells, have the ability to regenerate in culture without differentiation and have the ability to differentiate into specific mesenchymal lineages when either induced in vitro or in vivo at the site of damaged tissue. Human mesenchymal stem cells are known to differentiate under the appropriate conditions into multiple cell lineages, including osteoblasts, adipocytes and chondrocytes (for review see Caplan et al., 1997).

Accordingly, any process that is useful to recover mesenchymal stem cells from human tissue may be utilized to obtain a population of cells comprising enriched mesenchymal stem cells. In one aspect, the method of isolating human mesenchymal stem cells comprises the steps of providing a tissue specimen containing mesenchymal stem cells, preferably bone marrow; isolating the mesenchymal stem cells from the specimen, for example, by density centrifugation; adding the isolated cells to a medium which contains factors that stimulate mesenchymal stem cell growth without differentiation, and allows for the selective adherence of only the mesenchymal stem cells to a substrate surface in culture; culturing the specimen-medium mixture; and removing the non-adherent matter from the substrate surface.

In a further aspect of the present invention, any process that is useful to recover hematopoietic stem cells from human tissue may be utilized to result in a population of cells comprised mostly of hematopoietic cells. Stem cells can be recovered from various types of tissue such as bone marrow and blood, including peripheral blood. The human hematopoietic stem cells can be collected from bone marrow aspirates or peripheral blood and isolated using commercially available antibodies which bind to hematopoietic stem cell surface antigens, e.g. CD34, using methods known to those of skill in the art, see e.g. U.S. Pat. No. 4,714,680. Alternatively, the hematopoietic stem cells can be recovered by depletion, i.e., by selecting out cells which express markers found on other cells. The antibodies used for these procedures may be conjugated to magnetic beads and immunogenic procedures may be utilized to recover the desired cell type.

25

The human mesenchymal stem cells and the hematopoietic stem cells are co-cultured under appropriate culture conditions such that the mesenchymal stem cells adhere to a substrate surface and form a monolayer. The mesenchymal stem cells are plated at a density in a range of from about 3×10^3 to about 5×10^3 cells per cm^2 . The CD34+ cells are preferably at a cell density of approximately 5×10^4 cells per cm^2 .

30

According to the method of the present invention, the isolated mesenchymal stem cells and the isolated hematopoietic progenitor cells, preferably CD34+ cells, are

each culture expanded in appropriate media, i.e., cultured by methods using conditions that are apparent to those of skill in the art which favor cell growth and production of homogeneous cell populations.

5 The two cell populations are then co-cultured in a medium that promotes the growth of the human mesenchymal stem cells and does not adversely affect the maintenance of the hematopoietic stem cells. The media should also sustain the CD34+ cell population and support the induction of the differentiation of CD34+ cells into osteoclast cells. Suitable media are described for example in U.S. Pat. No.
10 5,486,359.

 In a preferred embodiment, the human mesenchymal stem cells and the hematopoietic stem cells are co-cultured in Dulbecco's Modified Eagles Medium-Low Glucose (DMEM-LG #11885 Life Technologies, Gaithersburg, MD). The medium
15 preferably contains 10% FBS. In a particularly preferred embodiment, the human mesenchymal stem cells and the hematopoietic stem cells are co-cultured in Dulbecco's Modified Eagles Medium-High Glucose (DMEM-HG # 11995 Life Technologies, Gaithersburg, MD) without FBS.

20 The hematopoietic stem cells produced according to the methods described herein can be used to provide a reliable and constant source of osteoclast cells. These osteoclasts could be used in the determination of the effects of osteotropic factors on the development and treatment of osteoporosis. It is contemplated that a ready supply of osteoclasts would be useful for screening assays for small molecule
25 drug development, e.g., based on expression of integrin $\alpha_v\beta_3$ cell surface receptor by osteoclasts. It is believed that these cells, which have not been treated with growth factors *in vitro*, will have characteristics more like osteoclasts found *in vivo* than have heretofore been obtained.

30 After differentiation of the hematopoietic progenitor cells into osteoclast cells as described herein, the mixture of hematopoietic progenitor cells, osteoclasts and mesenchymal stem cells may be separated to obtain a population of cells largely consisting of the osteoclast cells. This may be accomplished by positive

and/or negative selection of hematopoietic cells using antibodies to identify hematopoietic cell surface markers or other cell-type markers.

Another aspect of the present invention relates to the introduction of foreign
5 genes into the hematopoietic progenitor cells such that progeny of the cells, the osteoclasts, carry the new genetic material.

Thus, in accordance with this aspect of the invention, the hematopoietic
progenitor cells can be modified with genetic material of interest. The modified
10 CD34+ cells can then be co-cultured *in vitro* with mesenchymal stem cells and induced to differentiate into osteoclast cells. The osteoclast cells are able to express the product of the gene expression or secrete the expression product. These modified cells can then be administered to a target tissue, e.g. bone marrow, where the expressed product will have a beneficial effect.

15

In a further embodiment, the transduced hematopoietic progenitor cells can be induced *in vivo* to differentiate into osteoclasts that will express the gene product. For example, the transduced hematopoietic progenitor cells in combination with mesenchymal stem cells may be administered to induce
20 production of osteoclasts having the transduced gene. The cells may be administered in admixture with each other or separately and may be delivered to a targeted area. Alternatively, the hematopoietic progenitor cells may be used alone and caused to differentiate *in vivo* by mesenchymal stem cells present *in vivo*.

25 Thus, genes can be introduced into cells which are then returned to the autologous donor or an allogeneic recipient where the expression of the gene will have a therapeutic effect. For example, osteoclasts may be genetically engineered to have reduced activity *in vivo*. Appropriate genes would include those that play a role in the regulation of osteoporosis, in areas such as serum calcium
30 responsiveness, estrogen secretion and bone resorption.

The hematopoietic stem cells may be genetically modified (transduced or transformed or transfected) in the presence of the human mesenchymal stem cells, wherein the mesenchymal stem cells increase the efficiency of gene transduction of

the hematopoietic stem cells. Alternatively, the hematopoietic stem cells may be transduced in the absence of the human mesenchymal stem cells.

5 The hematopoietic stem cells may be genetically modified by incorporation of genetic material into the cells, for example using recombinant expression vectors.

As used herein "recombinant expression vector" refers to a transcriptional unit comprising an assembly of (1) a genetic element or elements having a regulatory role in gene expression, for example, promoters or enhancers, (2) a structural or coding
10 sequence which is transcribed into mRNA and translated into protein, and (3) appropriate transcription initiation and termination sequences. Structural units intended for use in eukaryotic expression systems preferably include a leader sequence enabling extracellular secretion of translated protein by a host cell. Alternatively, where recombinant protein is expressed without a leader or transport
15 sequence, it may include an N-terminal methionine residue. This residue may or may not be subsequently cleaved from the expressed recombinant protein to provide a final product.

The human hematopoietic progenitor cells thus may have stably integrated a
20 recombinant transcriptional unit into chromosomal DNA or carry the recombinant transcriptional unit as a component of a resident plasmid. Cells may be engineered with a polynucleotide (DNA or RNA) encoding a polypeptide *ex vivo*, for example. Cells may be engineered by procedures known in the art by use of a retroviral particle containing RNA encoding a polypeptide.

25 Retroviruses from which the retroviral plasmid vectors hereinabove mentioned may be derived include, but are not limited to, Moloney Murine Leukemia Virus, spleen necrosis virus, retroviruses such as Rous Sarcoma Virus, Harvey Sarcoma Virus, avian leukosis virus, gibbon ape leukemia virus, human
30 immunodeficiency virus, adenovirus, Myeloproliferative Sarcoma Virus, and mammary tumor virus. In one embodiment, the retroviral plasmid vector is MGIN, derived from murine embryonic stem cells. Generally regarding retroviral mediated gene transfer, see McLachlin et al.(1990).

The nucleic acid sequence encoding the polypeptide is under the control of a suitable promoter. Suitable promoters which may be employed include, but are not limited to, TRAP promoter, adenoviral promoters, such as the adenoviral major late promoter; the cytomegalovirus (CMV) promoter; the respiratory syncytial virus (RSV) promoter; inducible promoters, such as the MMT promoter, the metallothionein promoter; heat shock promoters; the albumin promoter; the ApoAI promoter; human globin promoters; viral thymidine kinase promoters, such as the Herpes Simplex thymidine kinase promoter; retroviral LTRs; ITRs; the β -actin promoter; and human growth hormone promoters. The promoter also may be the native promoter that controls the gene encoding the polypeptide. These vectors also make it possible to regulate the production of the polypeptide by the engineered progenitor cells. For example, for purposes of the present invention, the vector can contain a calcium response element. The selection of a suitable promoter will be apparent to those skilled in the art.

It is also possible to use vehicles other than retroviruses to genetically engineer or modify the hematopoietic stem cells. Genetic information of interest can be introduced by means of any virus which can express the new genetic material in such cells. For example, SV40, herpes virus, adenovirus and human papillomavirus can be used for this purpose. Other methods can also be used for introducing cloned eukaryotic DNAs into cultured mammalian cells, for example, the genetic material to be transferred to stem cells may be in the form of viral nucleic acids.

In addition, the expression vectors may contain one or more selectable marker genes to provide a phenotypic trait for selection of transformed cells such as dihydrofolate reductase or neomycin resistance.

The hematopoietic progenitor cells may be transfected through other means known in the art. Such means include, but are not limited to transfection mediated by calcium phosphate or DEAE-dextran; transfection mediated by the polycation Polybrene; protoplast fusion; electroporation; liposomes, either through encapsulation of DNA or RNA within liposomes, followed by fusion of the liposomes with the cell

membrane or, DNA coated with a synthetic cationic lipid can be introduced into cells by fusion.

The present invention further makes it possible to genetically engineer human hematopoietic progenitor cells in such a manner that they produce, *in vitro* or *in vivo* osteoclasts which produce polypeptides, hormones and proteins not normally produced in human osteoclasts in biologically significant amounts or produced in small amounts but in situations in which regulatory expression would lead to a therapeutic benefit. For example, the hematopoietic stem cells could be engineered with a gene that expresses a molecule that specifically inhibits bone resorption, but does not otherwise interfere with osteoclasts binding to bone. Alternatively the cells could be modified such that a protein normally expressed will be expressed at much lower levels. These products would then be secreted into the surrounding media or purified from the cells. The human osteoclast cells formed in this way can serve as continuous short term or long term production systems of the expressed substance.

This technology may be used to produce additional copies of essential genes to allow augmented expression by the osteoclasts of certain gene products *in vivo*. These genes can be, for example, hormones, matrix proteins, cell membrane proteins, cytokines, adhesion molecules, "rebuilding" proteins important in tissue repair. The expression of the exogenous genetic material *in vivo*, is often referred to as "gene therapy." Disease states and procedures for which such treatments have application include genetic disorders and diseases of bone and the immune system. Cell delivery of the transformed cells may be effected using various methods and includes infusion and direct depot injection into periosteal, bone marrow and subcutaneous sites.

In addition, as hereinabove described, the transduced cells may be used for *in vitro* production of desired protein(s). The transduced cells may further be used in screening assays for drug discovery.

In a preferred embodiment, hematopoietic stem cells are transduced with a retroviral vector containing a calcium response element, tartrate resistant acid phosphatase (TRAP) promoter and a cDNA encoding for anti-osteoclast factor.

After modification of the hematopoietic cells as described herein and induction of differentiation into osteoclasts, the mixture of hematopoietic, osteoclast and mesenchymal stem cells may be separated to obtain a population of cells largely consisting of the transduced osteoclasts. This may be accomplished
5 by positive and/or negative selection of transduced hematopoietic cells using antibodies to identify hematopoietic cell surface markers.

The above description of the invention and the following examples are by way of illustration only. Other permutations and practices of the invention will be
10 readily envisaged by one of ordinary skill in the art by view of the above in conjunction with the appended drawings. Therefore, such permutations and variations are within the scope of the present invention.

Examples

15

Statistical Analysis All data were analyzed by a paired t test. Samples were run in triplicates and data shown were means + SE.

Example 1

20

Cell isolation Bone marrow samples were collected from healthy human donors at the Johns Hopkins University under an Institutional Review Board-approved protocol. Mesenchymal stem cells (MSCs) were isolated and cultured according to known methods (Majumdar et al., 1998). When the cultures reached 90% confluence, adherent cells were recovered by the addition of a solution
25 containing 0.25% trypsin-EDTA (Life Technologies) and replated at a density of 1×10^6 cells per 185 cm^2 flask as passage-1 cells.

CD34+ cells, isolated from bone marrow of healthy patients were obtained from Poietic Technologies, Gaithersburg, MD). They were immunopurified to
30 95% purity using antibody to CD34 conjugated to magnetic beads (CD34+ cell separation column: Miltenyi Biotec, Auburn, CA Ab:QBEND.10) and cryopreserved.

Co-cultures MSCs of early passage (2 to 3) were plated at 3×10^3 cells per cm^2 and cultured until sub-confluency in DMEM LG+10% FBS. CD34⁺ cells were then added at 5×10^4 cells per cm^2 to MSC cultures. Cultures were maintained for 3 weeks at 37°C in an atmosphere of 95% air/5% CO₂ and fed every three days with the medium. Because most of the CD34⁺ cells remained non-adherent, half of the culture medium was gently aspirated without agitating non-adherent cells and replaced with fresh medium.

TRAP staining After three weeks, cells were fixed in 60% acetone in citrate buffer (pH 5.4) for 30 seconds, washed twice with distilled water, air-dried, and stained for tartrate-resistant acid phosphatase (TRAP), a widely-used cytochemical marker of mouse osteoclasts (Wijngaert and Burger, 1986), using a commercially available kit (Sigma, St. Louis, MO). Stained cultures were examined under light microscopy at a magnification of X200. TRAP-positive (red-staining) multinucleated (three or more nuclei) osteoclasts in each well were counted by manually scanning across the entire well in a systematic fashion (Mbalaviele et al., 1998).

CD34⁺ cells cultured in the absence of MSCs or cytokines/growth factors failed to form multinucleated cells expressing tartrate-resistant acid phosphatase (TRAP⁺MNCs) (Figure 1A). MSCs supported the differentiation of co-cultured CD34⁺ cells towards osteoclast cells (Figure 1B). In contrast, human skin fibroblasts (SK1087 cells) or human kidney cells (293T) failed to support the formation of TRAP⁺MNCs (Figures 1C and 1D, respectively). Treatment of CD34⁺ cells with IL-1, IL-3 and GM-CSF in the absence of MSCs yielded the formation of osteoclast cells, but the numbers of these cells were lower compared to those present in CD34⁺/MSCs co-cultures (data not shown).

Example 2

Immunocytochemistry CD34⁺ cells and MSCs co-cultured for three weeks were fixed with a fresh solution of 3.7% (wt./vol.) formaldehyde in PBS, blocked with 1.5% horse serum, incubated with the monoclonal antibody to human integrin $\alpha v \beta 3$ (Dr. Horton, The Rayne Institute, London, England) at 1:50 dilution in PBS

containing 0.15% horse serum. Cells were then incubated with glucose oxidase-conjugated goat secondary antibody using Vectastain ABC-GO kit (Vector Laboratories, Inc., Burlingame, CA). All incubations were performed at room temperature for 30 minutes followed by washing 3 times with PBS. Results are shown in Figure 2. Osteoclast cells (black arrow) and their precursors (arrowhead) were stained by vitronectin receptors antibody but not by the control antibody. MSCs did not show any staining with the receptor antibody. Multinucleated cells expressing low levels of vitronectin receptors were seen in co-cultures (Figure 2B, white arrow).

RNA Preparation and RT-PCR Total RNA was extracted from MSC cultures or co-cultures of CD34⁺ and MSCs using High Pure RNA Isolation kit purchased from Boehringer Mannheim (Indianapolis, IN). PCR was performed for 30 cycles either on reverse-transcribed single-strand cDNA from total RNA preparations (1 µg) using GeneAmp RT-PCR method under the conditions described by the supplier (Perkin Elmer, Foster City, CA) with the following modifications: denaturation at 95°C for 20 seconds, annealing at 55°C for 20 seconds, polymerization at 72°C for 30 seconds and elongation at 72°C for 10 minutes. The upstream and downstream primers, respectively, were designed as follows: TRAP: 5'-CGATCACAATCTGCAGTACC-3' (SEQ ID NO: 1) and 5'-ACCCAGTGAGTCTTCAGTCC-3' (SEQ ID NO:2), PCR product size=150 bp; calcitonin receptor (CTR): 5'-TTTCCAGGGCTTCTTTGTT-3' (SEQ ID NO:3) and 5'-CTTGGTTGTTGGCTGGTTC-3' (SEQ ID NO:4), PCR product size=205 bp. PCR products were size-fractionated by 1% agarose gel electrophoresis.

RT-PCR analysis showed expression of TRAP and calcitonin receptors by cells derived from CD34⁺ cells, the osteoclast lineage (Lanes 2 and 3 of Figure 3). These proteins were not expressed by MSCs.

Example 3

Pit formation assay To further confirm the presence of osteoclast cells in the cultures, the ability of cells produced in co-cultures of CD34⁺ cells and MSCs to form pits was examined, an indication of osteoclastic bone resorption. Elephant smoothened tusk slices were sterilized with 200-proof ethanol, air-dried and

washed several times with PBS. MSCs were plated at 3×10^3 cells per cm^2 on the tusk slices ($4 \times 4 \times 0.1$ mm) in 96-well plates and cultured for a week in hMSC medium pH 7.4. CD34^+ cells were then added at 5×10^4 cells per cm^2 . Artificial bone analog, osteologic discs (Millenium Biologix Inc., Ontario, Canada) were used to establish the optimal conditions of the assay. At the end of the 3-week culture period, the cells on slices were stained for TRAP, counted, then removed with 0.1 M NaOH in water followed by ultrasonication for 2 minutes. The bone slices were stained for 5 minutes with 1% toluidine blue prepared in distilled water containing 1% sodium borate. The numbers of pits were counted under light microscope as previously described (Prallet et al., 1992). Authentic resorption pits were found on bone analog as well as on elephant tusk slices when CD34^+ cells were co-cultured with MSCs (Figure 4).

Example 4

Effects of cell-cell interactions CD34^+ cells and MSCs were either cultured in the same compartment (contact co-cultures) or in two different compartments (non-contact co-cultures) separated by culture proof inserts with a $0.45 \mu\text{m}$ porous membrane (Becton Dickinson, Bedford, MA) to investigate physical cell-cell interactions. The results shown in Figure 5A indicated that osteoclast cells were formed in the contact cultures, and there were about two-thirds less osteoclast cells in non-contact cultures, regardless of in which compartment (top or bottom) the MSCs or CD34^+ cells were seeded.

Example 5

Interactions through secreted factors CD34^+ cells were co-cultured with MSCs. Conditioned media were collected after two days from MSCs cultured in the absence or presence of CD34^+ cells. The protein levels of cytokines IL- 1α , IL-6, IL-11, granulocyte/macrophage-colony stimulating factor (GM-CSF) and leukaemia inhibitory factor (LIF) secreted into the hematopoietic stem cell and mesenchymal stem cell co-cultures were measured using a Quantikine kit (R&D Systems, Minneapolis, MN).

Figure 5B shows the levels of IL-6, IL-11 or LIF in co-cultures of CD34⁺ cells and MSCs. The levels of IL-1 α and GM-CSF were undetectable. The levels of IL-6 and LIF were about 10 times greater in co-cultures of MSCs and CD34⁺ cells compared to MSCs cultures without CD34⁺ cells. The levels of IL-1 α and GM-CSF were undetectable.

Example 6

Effects of osteogenic MSCs (differentiated MSCs) Osteogenic differentiation of MSCs was induced by treating MSCs with osteogenic supplement (OS) containing 100mM dexamethasone 10mM β -glycerophosphate and 50 μ M L-ascorbic acid-2 phosphate for 2, 3, 4, 5, 10 or 13 days. At the time of the co-culture with CD34⁺ cells, OS containing medium was replaced with medium without OS. Figures 6A and 6B show that osteogenic MSCs inhibited TRAP⁺MNC formation compared to MSCs. Figure 6C shows the levels of IL-6, IL-11 or LIF in co-culture. The levels of IL-6, IL-11 and LIF were higher in co-cultures of MSCs with CD34⁺ cells compared to cocultures of osteogenic MSCs and CD34⁺ cells.

RNA preparation and RT-PCR Total RNA isolation and RT-PCR analysis were carried out as described in example 2. The upstream and downstream primers, respectively, for OPG were 5'-ACCACTACTACACAGACAGC-3' (SEQ ID NO:5) and 5'-AGGAGACCAAAGACACTGCA-3' (SEQ ID NO:6) ; for OPGL 5'-TTCTATTTTCAGAGCGCAGAT-3' (SEQ ID NO:7) and 5'-AGTCATGTTGGAGATCTTGG-3' (SEQ ID NO:8). Figure 6D shows that treatment of MSCs with OS for 4, 8, or 15 days decreased the expression of OPGL mRNA. In contrast, treatment of MSCs with OS for 4, 8 or 15 days increased the expression of OPG mRNA.

Example 7

Transduction of CD34⁺ cells The construction of the retroviral vector MGIN expressing enhanced green fluorescent protein (EGFP) gene and the production of the retroviral supernatants has been previously reported (Cheng et al., 1997). Briefly, MGIN is a murine embryonic stem cell virus-based vector

containing EGFP gene and the internal ribosome entry site (IRES). Amphotropic supernatants produced by PA317 packaging cells were made from selected producers after infection by BOS23 ecotropic viral stocks. For transduction, previously frozen vector supernatants were mixed at a 1:1 ratio with medium containing CD34⁺ cells in the presence of 8 µg/ml polybrene (Sigma, St. Louis, MO), interleukin-3 (IL-3) and IL-6 (10 ng/ml, each), stem cell factor (SCF) and Flk2 (FL) (100 ng/ml, each). Control cells were transduced with non-EGFP-expressing vector. The transduction mixture was then centrifuged at 1800 g at 32-35°C. After a 4-hour "spinoculation," cells were washed once and cultured for 24 hours in medium containing the above growth factors. After the transduction was repeated once more, a cell fraction was analyzed by flow cytometry to test transduction efficiency while the remaining cells were used for the osteoclastogenesis assay. The transduced CD34⁺ cells were co-cultured for three weeks without added factors. The co-cultures were stained for TRAP as described in example 1. Cells were visualized with a fluorescence microscope to detect EGFP or under a light microscope to detect TRAP.

Flow cytometry analysis results are shown in Figure 7. FACS analysis showed that 30% of CD34⁺ cells transduced with EGFP vector expressed the transgene (Figure 7 B). Co-cultures of transduced CD34⁺ cells with MSCs yielded osteoclasts expressing both EGFP (Figure 7C) and TRAP (Figure 7D).

Example 8

Detection of vitronectin receptor by ELISA MSCs were plated at 3×10^3 cells per cm² and cultured in hMSC medium until subconfluency. CD34⁺ cells were then added at 2.5×10^4 cells per cm² to MSC cultures. Cultures were maintained for 1, 2 or 3 weeks. Cells were fixed with 3.6% formaldehyde and blocked with 10% goat serum for 30 minutes, followed by incubation with the antibody to the vitronectin receptor ($\alpha v \beta 3$). With extensive washes between additions, a biotin-labeled secondary antibody was added (goat anti-mouse IgG; Pierce, Rockford, IL) followed by streptavidin-conjugated β -galactosidase (Gibco BRL) for 1 hour. The soluble substrate chlorophenol red- β -D-galactopyranoside (CPRG; Boehringer Mannheim, Indianapolis, IN) was then added, and color

development was read at 570 nm after 30 minutes. Control primary antibodies were also run for each treatment and this background measurement was subtracted from the readings for each well. ELISA results are presented in Figure 8 as mean absorbance units, with background subtracted. The vitronectin receptor ($\alpha v\beta 3$) was detected by the specific antibody but not by the control antibody (Figure 8A). $\alpha v\beta 3$ expression by the osteoclast cell lineage was induced after 1 week of co-cultures and was less expressed by MSCs (Figure 8B).

Example 9

Hematopoietic stem cells as therapeutic targets Hematopoietic stem cells (HSCs) are transduced with a retroviral vector containing a calcium response element, tartrate resistant acid phosphatase (TRAP) promoter and a cDNA encoding for anti-osteoclast factor (antisense, inhibitor, apoptotic factor) (see Figure 8). TRAP promoter has been used to specifically target the osteoclast cell lineage. Transduced-HSCs are infused to osteoporotic patients. They home to bone marrow, engraft and differentiate into osteoclasts. The osteoclasts respond to local high calcium concentrations generated by excessive bone resorption (steps 1 and 2), by expressing an anti-osteoclast factor (step 3) which stops osteoclast activity (step 4). One such factor is Osteoprotegerin (OPG) which when secreted has been demonstrated to decrease bone resorption (osteopetrosis) and appears to generally block osteoclastogenesis (Simonet (1997)). The transduced hematopoietic cells can also be induced to differentiate into osteoclasts ex vivo, and then infused to osteoporotic patients.

Calcium-response element (CaRE) was cloned under the control of CMV promoter within pGL3 vector that contains luciferase as a reporter gene (Promega Madison WI) to yield pGL3-CaRE. pGL3 or pGL3-CaRE were transfected into human kidney cell line 293T using a calcium phosphate co-precipitation method. Cells were then cultured in the presence of increasing concentrations of calcium. Luciferase activity was assessed using Dual-Luciferase™ reporter assay system (Promega). The data show that calcium as low as 0.1mM induced luciferase activity in pGL3-CaRE but not in pGL3 (Figure 9B).

Example 10

Effect of serum in culture medium Co-cultures of MSCs and CD34+ cells were studied to determine the effect of serum on the production of osteoclasts. Co-cultures of CD34+ cells and MSCs were cultured, under the conditions described above, however the media tested were DMEM-low glucose with 10% FBS and DMEM-high glucose without serum. Control cultures contained human kidney cell line 293 cells co-cultured with CD34+ cells in DMEM-high glucose with and without serum. The cultures were terminated at 12 days rather than 21 days as in the experiments in the foregoing examples.

10

The results shown in Figure 10 demonstrate that co-culturing MSCs and CD34+ cells in DMEM-high glucose without FBS yielded a greater number of osteoclasts than co-culture in the standard DMEM-LG with 10% FBS.

CITED LITERATURE

Caplan AI, Bruder SP: Cell and molecular engineering of bone regeneration, in Lanza R, Langer R, Chick W (eds): Textbook of Tissue Engineering, Georgetown, Landes Company, 1997, p 603

5

Cheng L, Du C, Murray D, Tong X, Zhang YA, Chen BP, Hawley RG: A GFP reporter system to assess gene transfer and expression in human hematopoietic progenitor cells. *Gene Therapy* 4: 1013 (1997)

- 10 James IE, Dodds RA, Lee-Rykaczewski E, Eichman CF, Connor JR, Hart T, Maleef BE, Lackman RD, Gowen M: Purification and characterization of fully functional human osteoclast precursors. *J. Bone Min. Res.* 11: 1608 (1996)

- 15 Kong YY, Yoshida H, Sarosi I, Tan HL, Timms E, Capparelli C, Morony S, Oliveira-dos-Santos AJ, Van G, Itie A, Khoo W, Wakeham A, Dunstan CR, Lacey DL, Mak TW, Boyle WJ, Penninger JM: OPGL is a key regulator of osteoclastogenesis, lymphocyte development and lymph-node organogenesis. *Nature* 397(6717):315-23 (1999)

- 20 Lacey DL, Timms E, Tan HL, Kelley MJ, Dunstan CR, Burgess T, Elliott R, Colombero A, Elliott G, Scully S, Hsu H, Sullivan J, Hawkins N, Davy E, Capparelli C, Eli A, Qian YX, Kaufman S, Sarosi I, Shalhoub V, Senaldi G, Guo J, Delaney J, Boyle WJ: Osteoprotegerin ligand is a cytokine that regulates osteoclast differentiation and activation. *Cell* 93(2):165-76 (1998)

25

McLachlin et al. Progress in Nucleic Acid Research and Molecular Biology, 38:91-135(1990)

- 30 Majumdar MK, Thiede MA, Mosca JD, Moorman M, Gerson S: Phenotypic and functional comparison of cultures of marrow-derived mesenchymal stem cells (MSCs) and stromal cells. *J Cell. Physiol.* 176(1):57-66 (1998)

Matayoshi A, Brown C, DiPersio JF, Haug J, Abu-Amer Y, Liapis H, Kuestner R, Pacifici R: Human blood-mobilized hematopoietic precursors differentiate into

osteoclasts in the absence of stromal cells. *Proc. Natl. Acad. Sci. USA* 93, 10785 (1996)

5 Mbalaviele G, Orcel P, Morieux C, Nijweide PJ, Devemejoul MC: Osteoclast formation from human cord blood mononuclear cells co-cultured with mice embryonic metatarsals in the presence of M-CSF. *Bone* 16: 171 (1995)

10 Mbalaviele G, Nishimura R, Myoi A, Niewolna M, Reddy SV, Chen D, Feng J, Roodman D, Mundy GR, Yoneda T: Cadherin-6 mediates the heterotypic interactions between the hemopoietic osteoclast cell lineage and stromal cells in a murine model of osteoclast differentiation. *J. Cell Biol* 141(6):1467-1476 (1998).

15 Mundy GR, Boyce BF, Yoneda T, Bonewald LF, Roodman GD: Cytokines and bone remodeling, in Marcus R, Feldman D, Kelsey J (eds): Osteoporosis, San Diego, Academic Press, 1995, p 302

20 Prallet B, Male P, Neff L, Baron R: Identification of a functional mononuclear precursor of the osteoclast in chicken medullary bone marrow cultures. *J Bone Min. Res.* 7: 405 (1992)

Sarma U, Flanagan AM: Macrophage colony-stimulating factor induces substantial osteoclast generation and bone resorption in human bone marrow cultures. *Blood*, 88: 2531 (1996)

25 Quinn JMW, Fujikawa Y, McGee JO'D, Athanasou NA: Rodent osteoblast-like cells support osteoclastic differentiation of human cord blood monocytes in the presence of M-CSF and 1,25 Dihydroxyvitamin D3. *Int. J. Biochem. Cell Biol.* 29: 173 (1997)

30 Simonet et al. Osteoprotegerin: A Novel Secreted Protein Involved in the Regulation of Bone Density. *J. Biol. Chem.* 268(1):377-384 (1993)

Takahashi S, Reddy SV, Dallas M, Devlin R, Chou JY, Roodman GD: Development and characterization of a human marrow stromal cell line that enhances osteoclast-like cell formation. *Endocrinol.* 136: 1441 (1995)

- 5 Takahashi S. et al. *Endocrinology* 123:2600-2602 (1988)

Teitelbaum SL, Tondravi MM, Ross FP: Osteoclast biology, in Marcus R, Feldman D, Kelsey J (eds): Osteoporosis, San Diego, Academic Press, 1995, p 61

- 10 Wijngaert FB, Burger EH: Demonstration of tartrate-resistant acid phosphatase in undecalcified, glycomethacrylate-embedded mouse bone: a possible marker for (pre-osteoclast identification *J Histol. Cytol.* 34:1317 (1986)

CLAIMS

1. A method for producing osteoclasts in vitro comprising co-culturing
5 mesenchymal stem cells with hematopoietic stem cells such that the hematopoietic stem cells produce osteoclasts.
2. The method of claim 2 wherein co-culturing is done in the absence of
exogenous cytokines.
- 10 3. The method of claim 1 wherein the hematopoietic stem cells are CD34+ cells.
4. The method of claim 1 wherein the hematopoietic stem cells and the
15 mesenchymal stem cells are in contact.
5. A method of producing genetically modified osteoclasts, comprising
transducing hematopoietic progenitor cells with exogenous genetic material; and
culturing the transduced hematopoietic cells in the presence of mesenchymal stem
20 cells to induce differentiation of the transduced hematopoietic cells into osteoclasts which contain the exogenous genetic material.
6. The method of claim 5 wherein the transduced hematopoietic stem cells
are induced to differentiate *in vitro*.
- 25 7. The method of claim 5 wherein the transduced hematopoietic stem cells are induced to differentiate *in vivo*.
8. A method of obtaining genetically modified osteoclasts comprising
30 transducing hematopoietic progenitor cells with exogenous genetic material and placing the transduced hematopoietic cells under conditions suitable for differentiation of the hematopoietic stem cells into osteoclast cells which contain the exogenous genetic material.

9. The method of claim 8 wherein the hematopoietic cells are differentiated *in vivo*.

10. The method of claim 8 wherein the hematopoietic cells are administered *in vivo* with human mesenchymal stem cells.

11. A composition comprising genetically modified osteoclasts.

12. The composition of claim 11 wherein the genetically modified osteoclasts are from a non-transformed cell line.

13. A composition comprising genetically modified osteoclast precursor cells.

14. The composition of claim 13 wherein the genetically modified osteoclast precursor cells are from a non-transformed cell line.

15. A method for decreasing differentiation of hematopoietic stem cells into osteoclasts comprising co-culturing differentiating mesenchymal stem cells with hematopoietic stem cells such that osteoclastogenesis is reduced.




FIG. 1A

FIG. 1B





FIG. 1C

FIG. 1D




FIG. 2A





FIG. 2B



FIG. 3



+

FIG. 4A






FIG. 4B

Pit number/osteoclast-like cell

FIG. 4C



+

FIG. 5A




FIG. 5B




FIG. 6A




FIG. 6B




FIG. 6C





FIG. 6D






FIG. 7B





FIG. 7A




FIG. 7C

FIG. 7D

FIG. 8A




FIG. 8B




FIG. 9B




FIG. 9A




FIG. 10B





FIG. 10A



SEQUENCE LISTING

<110> Osiris Therapeutics, Inc.

<120> Regulation of Hematopoietic Stem Cell Differentiation
by the Use of Human Mesenchymal Stem Cells

<130> 640100-333

<140>

<141>

<150> 60/099,233

<151> 1998-09-04

<150> 60/088,431

<151> 1998-06-08

<160> 8

<170> PatentIn Ver. 2.0

<210> 1

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: PCR Primer

<400> 1

cgatcacaat ctgcagtacc

20

<210> 2

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: PCR Primer

<400> 2

accagtgag tcttcagtcc

20

<210> 3

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: PCR Primer

<400> 3

tttccagggc ttctttgtt

19

<210> 4

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: PCR Primer

<400> 4

cttggttggt ggctgggtc

19

<210> 5

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: PCR Primer

<400> 5

accactacta cacagacagc

20

<210> 6

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: PCR Primer

<400> 6

aggagaccaa agacactgca

20

<210> 7

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: PCR Primer

<400> 7
ttctatttca gagcgagat 20

<210> 8
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: PCR Primer

<400> 8
agtcagtgtg gagatcttgg 20

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 99/12847

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C12N5/06 C12N5/08 C12N5/10 A61K35/12 A61K48/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C12N A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 670 351 A (EMERSON STEPHEN G ET AL) 23 September 1997 (1997-09-23) column 10, line 51 -column 11, line 67 claims 1,18,50,108 ---	8,11-14
A	WO 95 02040 A (GSF FORSCHUNGSZENTRUM UMWELT ;THALMEIER KARIN (DE); DOERMER PETER) 19 January 1995 (1995-01-19) the whole document ---	1-15
A	WO 96 32469 A (RODAN SEVGI B ;MERCK & CO INC (US); RODAN GIDEON A (US); WESOLOWSK) 17 October 1996 (1996-10-17) the whole document ---	1-15
A	US 5 486 359 A (CAPLAN ARNOLD I ET AL) 23 January 1996 (1996-01-23) the whole document ---	1-15
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance
"E" earlier document but published on or after the international filing date
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
"O" document referring to an oral disclosure, use, exhibition or other means
"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
"&" document member of the same patent family

Date of the actual completion of the international search

14 December 1999

Date of mailing of the international search report

20/12/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Fernandez y Branas, F

INTERNATIONAL SEARCH REPORT

Application No

PCT/US 95/12847

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>TAKAHASHI S ET AL: "Development and characterization of a human marrow stromal cell line that enhances osteoclast -like cell formation." ENDOCRINOLOGY, (1995 APR) 136 (4) 1441-9. , XP000857459 cited in the application page 1443, column 1, paragraph 3 page 1445, column 2, paragraphs 1,2</p>	1-15
T	<p>MBALAVIELE G ET AL: "Human mesenchymal stem cells promote human osteoclast differentiation from CD34+ bone marrow hematopoietic progenitors." ENDOCRINOLOGY, (1999 AUG) 140 (8) 3736-43. , XP000857472 the whole document</p>	1-15

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 99/12847

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 5 (partially), 7, 8 (partially), 9-10, 15 (partially) are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

information to next family members

PCT/U/12847

Form PCT/ISA/210 (patent family annex) (July 1992)

1/

2/